

LOGISTIC EQUATION WITH THE p -LAPLACIAN AND CONSTANT YIELD HARVESTING

SHOBHA ORUGANTI, JUNPING SHI,
AND RATNASINGHAM SHIVAJI

Received 29 September 2003

We consider the positive solutions of a quasilinear elliptic equation with p -Laplacian, logistic-type growth rate function, and a constant yield harvesting. We use sub-super-solution methods to prove the existence of a maximal positive solution when the harvesting rate is under a certain positive constant.

1. Introduction

We consider weak solutions to the boundary value problem

$$\begin{aligned} -\Delta_p u &= f(x, u) \equiv au^{p-1} - u^{y-1} - ch(x) && \text{in } \Omega, \\ u &> 0 && \text{in } \Omega, \\ u &= 0 && \text{on } \partial\Omega, \end{aligned} \tag{1.1}$$

where Δ_p denotes the p -Laplacian operator defined by $\Delta_p z := \operatorname{div}(|\nabla z|^{p-2} \nabla z)$; $p > 1$, $y (> p)$, a and c are positive constants, Ω is a bounded domain in R^N ; $N \geq 1$, with $\partial\Omega$ of class $C^{1,\beta}$ for some $\beta \in (0, 1)$ and connected (if $N = 1$, we assume Ω is a bounded open interval), and $h: \bar{\Omega} \rightarrow R$ is a continuous function in $\bar{\Omega}$ satisfying $h(x) \geq 0$ for $x \in \Omega$, $h(x) \not\equiv 0$, $\max_{x \in \bar{\Omega}} h(x) = 1$, and $h(x) = 0$ for $x \in \partial\Omega$. By a weak solution of (1.1), we mean a function $u \in W_0^{1,p}(\Omega)$ that satisfies

$$\int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla w \, dx = \int_{\Omega} [au^{p-1} - u^{y-1} - ch(x)] w \, dx, \quad \forall w \in C_0^\infty(\Omega). \tag{1.2}$$

From the standard regularity results of (1.1), the weak solutions belong to the function class $C^{1,\alpha}(\bar{\Omega})$ for some $\alpha \in (0, 1)$ (see [4, pages 115–116] and the references therein).

We first note that if $a \leq \lambda_1$, where λ_1 is the first eigenvalue of $-\Delta_p$ with Dirichlet boundary conditions, then (1.1) has no positive solutions. This follows since if u is a positive solution of (1.1), then u satisfies

$$\int_{\Omega} |\nabla u|^p \, dx = \int_{\Omega} [au^{p-1} - u^{y-1} - ch(x)] u \, dx. \tag{1.3}$$

But $\int_{\Omega} |\nabla u|^p dx \geq \lambda_1 \int_{\Omega} u^p dx$. Combining, we obtain $\int_{\Omega} [au^{p-1} - u^{y-1} - ch(x)]u dx \geq \lambda_1 \int_{\Omega} u^p dx$ and hence $\int_{\Omega} (a - \lambda_1)u^p dx \geq \int_{\Omega} [u^{y-1} + ch(x)]u dx \geq 0$. This clearly requires $a > \lambda_1$.

Next if $a > \lambda_1$ and c is very large, then again it can be proven that there are no positive solutions. This follows easily from the fact that if the solution u is positive, then $\int_{\Omega} [au^{p-1} - u^{y-1} - ch(x)]dx$ is nonnegative. In fact, from the divergence theorem (see [4, page 151]),

$$\int_{\Omega} [au^{p-1} - u^{y-1} - ch(x)]dx = - \int_{\partial\Omega} |\nabla u|^{p-2} \nabla u \cdot \nu dx \geq 0. \tag{1.4}$$

Thus,

$$c \int_{\Omega} h(x)dx \leq \int_{\Omega} [au^{p-1} - u^{y-1}]dx \leq a^{(y-1)/(y-p)} |\Omega|. \tag{1.5}$$

Here in the last inequality, we used the fact that $u(x) \leq a^{1/(y-p)}$ which can be proven by the maximum principle (see [4, page 173]).

This leaves us with the analysis of the case $a > \lambda_1$ and c small which is the focus of the paper.

THEOREM 1.1. *Suppose that $a > \lambda_1$. Then there exists $c_0(a) > 0$ such that if $0 < c < c_0$, then (1.1) has a positive $C^{1,\alpha}(\bar{\Omega})$ solution u . Further, this solution u is such that $u(x) \geq (ch(x)/\lambda_1)^{1/(p-1)}$ for $x \in \bar{\Omega}$.*

THEOREM 1.2. *Suppose that $a > \lambda_1$. Then there exists $c_1(a) \geq c_0$ such that for $0 < c < c_1$, (1.1) has a maximal positive solution, and for $c > c_1$, (1.1) has no positive solutions.*

Remark 1.3. **Theorem 1.2** holds even when $h(x) > 0$ in $\bar{\Omega}$.

We establish **Theorem 1.1** by the method of sub-supersolutions. By a supersolution (subsolution) ϕ of (1.1), we mean a function $\phi \in W_0^{1,p}(\Omega)$ such that $\phi = 0$ on $\partial\Omega$ and

$$\int_{\Omega} |\nabla \phi|^{p-2} \nabla \phi \cdot \nabla w dx \geq (\leq) \int_{\Omega} [a\phi^{p-1} - \phi^{y-1} - ch(x)]w dx, \quad \forall w \in W, \tag{1.6}$$

where $W = \{v \in C_0^\infty(\Omega) \mid v \geq 0 \text{ in } \Omega\}$. Now if there exist subsolutions and supersolutions ψ and ϕ , respectively, such that $0 \leq \psi \leq \phi$ in Ω , then (1.1) has a positive solution $u \in W_0^{1,p}(\Omega)$ such that $\psi \leq u \leq \phi$. This follows from a result in [3].

Equation (1.1) arises in the studies of population biology of one species with u representing the concentration of the species and $ch(x)$ representing the rate of harvesting. The case when $p = 2$ (the Laplacian operator) and $\gamma = 3$ has been studied in [6]. The purpose of this paper is to extend some of this study to the p -Laplacian case. In [3], the authors studied (1.1) in the case when $c = 0$ (nonharvesting case). However, the $c > 0$ case is a semipositone problem ($f(x,0) < 0$) and studying positive solutions in this case is significantly harder. Very few results exist on semipositone problems involving the p -Laplacian operator (see [1, 2]), and these deal with only radial positive solutions with the domain Ω a ball or an annulus. In **Section 2**, when $a > \lambda_1$ and c is sufficiently small, we will construct nonnegative subsolutions and supersolutions ψ and ϕ , respectively, such that $\psi \leq \phi$, and

establish [Theorem 1.1](#). We also establish [Theorem 1.2](#) in [Section 2](#) and discuss the case when $h(x) > 0$ in $\bar{\Omega}$.

2. Proofs of theorems

Proof of [Theorem 1.1](#). We first construct the subsolution ψ . We recall the antimaximum principle (see [[4](#), pages 155–156]) in the following form. Let λ_1 be the principal eigenvalue of $-\Delta_p$ with Dirichlet boundary conditions. Then there exists a $\delta(\Omega) > 0$ such that the solution z_λ of

$$\begin{aligned} -\Delta_p z - \lambda z^{p-1} &= -1 \quad \text{in } \Omega, \\ z &= 0 \quad \text{on } \partial\Omega, \end{aligned} \tag{2.1}$$

for $\lambda \in (\lambda_1, \lambda_1 + \delta)$ is positive for $x \in \Omega$ and is such that $(\partial z_\lambda / \partial \nu)(x) < 0, x \in \partial\Omega$.

We construct the subsolution ψ of [\(1.1\)](#) using z_λ such that $\lambda_1 \psi(x)^{p-1} \geq ch(x)$. Fix $\lambda_* \in (\lambda_1, \min\{a, \lambda_1 + \delta\})$. Let $\alpha = \|z_{\lambda_*}\|_\infty, K_0 = \inf\{K \mid \lambda_1 K^{p-1} z_{\lambda_*}^{p-1} \geq h(x)\}$, and $K_1 = \max\{1, K_0\}$. Define $\psi = K c^{1/(p-1)} z_{\lambda_*}$, where $K \geq K_1$ is to be chosen. Let $w \in W$. Then

$$\begin{aligned} & - \int_{\Omega} |\nabla \psi|^{p-2} \nabla \psi \cdot \nabla w \, dx + \int_{\Omega} [a(\psi)^{p-1} - (\psi)^{y-1} - ch(x)] w \, dx \\ &= \int_{\Omega} [-cK^{p-1}(\lambda_* z_{\lambda_*}^{p-1} - 1) + ac(Kz_{\lambda_*})^{p-1} - (Kc^{1/(p-1)} z_{\lambda_*})^{y-1} - ch(x)] w \, dx \\ &\geq \int_{\Omega} [-cK^{p-1}(\lambda_* z_{\lambda_*}^{p-1} - 1) + ac(Kz_{\lambda_*})^{p-1} - (Kc^{1/(p-1)} z_{\lambda_*})^{y-1} - c] w \, dx \\ &= \int_{\Omega} [(a - \lambda_*)(Kz_{\lambda_*})^{p-1} - (Kz_{\lambda_*})^{y-1} c^{(y-p)/(p-1)} + (K^{p-1} - 1)] c w \, dx. \end{aligned} \tag{2.2}$$

Define $H(y) = (a - \lambda_*)y^{p-1} - y^{y-1}c^{(y-p)/(p-1)} + (K^{p-1} - 1)$. Then $\psi(x)$ is a subsolution if $H(y) \geq 0$ for all $y \in [0, K\alpha]$. But $H(0) = K^{p-1} - 1 \geq 0$ since $K \geq 1$ and $H'(y) = y^{p-2}[(a - \lambda_*)(p-1) - c^{(y-p)/(p-1)}(y-1)y^{y-p}]$. Hence $H(y) \geq 0$ for all $y \in [0, K\alpha]$ if $H(K\alpha) = (a - \lambda_*)(K\alpha)^{p-1} - (K\alpha)^{y-1}c^{(y-p)/(p-1)} + (K^{p-1} - 1) \geq 0$, that is, if

$$c \leq \left(\frac{(a - \lambda_*)(K\alpha)^{p-1} + (K^{p-1} - 1)}{(K\alpha)^{y-1}} \right)^{(p-1)/(y-p)}. \tag{2.3}$$

We define

$$c_1 = \sup_{K \geq K_1} \left(\frac{(a - \lambda_*)(K\alpha)^{p-1} + (K^{p-1} - 1)}{(K\alpha)^{y-1}} \right)^{(p-1)/(y-p)}. \tag{2.4}$$

Then for $0 < c < c_1$, there exists $\bar{K} \geq K_1$ such that

$$c < \left(\frac{(a - \lambda_*)(\bar{K}\alpha)^{p-1} + (\bar{K}^{p-1} - 1)}{(\bar{K}\alpha)^{y-1}} \right)^{(p-1)/(y-p)} \tag{2.5}$$

and hence $\psi(x) = \bar{K} c^{1/(p-1)} z_{\lambda_*}$ is a subsolution.

We next construct the supersolution $\phi(x)$ such that $\phi(x) \geq \psi(x)$. Let $G(y) = ay^{p-1} - y^{y-1}$. Since $G'(y) = y^{p-2}[a(p-1) - (y-1)y^{y-p}]$, $G(y) \leq L = G(y_0)$, where $y_0 = [a(p-1)/(y-1)]^{1/(y-p)}$. Let ϕ be the positive solution of

$$-\Delta_p \phi = L \quad \text{in } \Omega, \tag{2.6}$$

$$\phi = 0 \quad \text{on } \partial\Omega. \tag{2.7}$$

Then for $w \in W$,

$$\begin{aligned} \int_{\Omega} |\nabla \phi|^{p-2} \nabla \phi \cdot \nabla w \, dx &= \int_{\Omega} Lw \, dx \\ &\geq \int_{\Omega} [a\phi^{p-1} - \phi^{y-1}]w \, dx \\ &\geq \int_{\Omega} [a\phi^{p-1} - \phi^{y-1} - ch(x)]w \, dx. \end{aligned} \tag{2.8}$$

Thus ϕ is a supersolution of (1.1). Also since $-\Delta_p \psi \leq a\psi^{p-1} - \psi^{y-1} - ch(x) \leq L = -\Delta_p \phi$, by the weak comparison principle (see [4, 5]), we obtain $\phi \geq \psi \geq 0$. Hence there exists a solution $u \in W_0^{1,p}(\Omega)$ such that $\phi \geq u \geq \psi$. From the regularity results (see [4, pages 115–116] and the references therein), $u \in C^{1,\alpha}(\bar{\Omega})$. \square

Remark 2.1. If \tilde{u} is any $C^{1,\alpha}(\bar{\Omega})$ solution of (1.1), then by the weak comparison principle, $\|\tilde{u}\|_{\infty} \leq \|\phi\|_{\infty}$, where ϕ is as in (2.6).

Proof of Theorem 1.2. From Theorem 1.1, we know that for c small, there exists a positive solution. Whenever (1.1) has a positive solution u , (1.1) also has a maximal positive solution. This easily follows since ϕ in (2.6) is always a supersolution such that $\phi \geq u$. Next if for $c = \bar{c}$, we have a positive solution $u_{\bar{c}}$, then for all $c < \bar{c}$, $u_{\bar{c}}$ is a positive subsolution. Hence again using ϕ in (2.6) as the supersolution, we obtain a maximal positive solution for c . From (1.3), it is easy to see that for large c , there does not exist any positive solution. Hence there exists a $c_1(a) > 0$ such that there exists a maximal positive solution for $c \in (0, c_1)$ and no positive solution for $c > c_1$. \square

Remark 2.2. The use of the antimaximum principle in the creation of the subsolution helps us to easily modify the proof of Theorem 1.1 to obtain a positive maximal solution for all $c < c_2(a) = \sup_{K \geq 1} (((a - \lambda_*)(K\alpha)^{p-1} + (K^{p-1} - 1))/(K\alpha)^{y-1})^{(p-1)/(y-p)}$ even in the case when $h(x) > 0$ in $\bar{\Omega}$. Here $c_2(a) \geq c_0(a)$. (Of course when $h(x) > 0$ in $\bar{\Omega}$, our solution does not satisfy $u(x) \geq (ch(x)/\lambda_1)^{1/(p-1)}$ for $x \in \bar{\Omega}$.) Hence Theorem 1.2 also holds in the case when $h(x) > 0$ in $\bar{\Omega}$.

Acknowledgment

The research of the second author is partially supported by National Science Foundation (NSF) Grant DMS-0314736, College of William and Mary summer research Grants, and a Grant from Science Council of Heilongjiang Province, China.

References

- [1] H. Dang, K. Schmitt, and R. Shivaji, *On the number of solutions of boundary value problems involving the p -Laplacian*, Electron. J. Differential Equations **1996** (1996), no. 1, 1–9.
- [2] H. Dang, R. Shivaji, and C. Maya, *An existence result for a class of superlinear p -Laplacian semipositone systems*, Differential Integral Equations **14** (2001), no. 2, 231–240.
- [3] P. Drábek and J. Hernández, *Existence and uniqueness of positive solutions for some quasilinear elliptic problems*, Nonlinear Anal. Ser. A: Theory Methods **44** (2001), no. 2, 189–204.
- [4] P. Drábek, P. Krejčí, and P. Takáč, *Nonlinear Differential Equations*, Chapman & Hall/CRC Research Notes in Mathematics, vol. 404, Chapman & Hall/CRC, Florida, 1999.
- [5] J. Fleckinger-Pellé and P. Takáč, *Uniqueness of positive solutions for nonlinear cooperative systems with the p -Laplacian*, Indiana Univ. Math. J. **43** (1994), no. 4, 1227–1253.
- [6] S. Oruganti, J. Shi, and R. Shivaji, *Diffusive logistic equation with constant yield harvesting. I. Steady states*, Trans. Amer. Math. Soc. **354** (2002), no. 9, 3601–3619.

Shobha Oruganti: Department of Mathematics, School of Science, Penn State Erie, The Behrend College, Penn State University, PA 16563, USA

E-mail address: sxo12@psu.edu

Junping Shi: Department of Mathematics, College of William and Mary, Williamsburg, VA 23187-8795, USA; School of Mathematics, Harbin Normal University, Harbin, Heilongjiang 150080, China


E-mail address: shij@math.wm.edu

Ratnasingham Shivaji: Department of Mathematics and Statistics, Mississippi State University, Mississippi State, MS 39762, USA

E-mail address: shivaji@ra.msstate.edu

DISCRETE OSCILLATION THEORY

Ravi P. Agarwal, Martin Bohner, Said R. Grace, and Donal O'Regan



This book is devoted to a rapidly developing branch of the qualitative theory of difference equations with or without delays. It presents the theory of oscillation of difference equations, exhibiting classical as well as very recent results in that area. While there are several books on difference equations and also on oscillation theory for ordinary differential equations, there is until now no book devoted solely to oscillation theory for difference equations. This book is filling the gap, and it can easily be used as an encyclopedia and reference tool for discrete oscillation theory.

In nine chapters, the book covers a wide range of subjects, including oscillation theory for second-order linear difference equations, systems of difference equations, half-linear difference equations, nonlinear difference equations, neutral difference equations, delay difference equations, and differential equations with piecewise constant arguments. This book summarizes almost 300 recent research papers and hence covers all aspects of discrete oscillation theory that have been discussed in recent journal articles. The presented

theory is illustrated with 121 examples throughout the book. Each chapter concludes with a section that is devoted to notes and bibliographical and historical remarks.

The book is addressed to a wide audience of specialists such as mathematicians, engineers, biologists, and physicists. Besides serving as a reference tool for researchers in difference equations, this book can also be easily used as a textbook for undergraduate or graduate classes. It is written at a level easy to understand for college students who have had courses in calculus.

For more information and online orders please visit <http://www.hindawi.com/books/cmia/volume-1>
For any inquires on how to order this title please contact books.orders@hindawi.com

"Contemporary Mathematics and Its Applications" is a book series of monographs, textbooks, and edited volumes in all areas of pure and applied mathematics. Authors and/or editors should send their proposals to the Series Editors directly. For general information about the series, please contact cmia.ed@hindawi.com.