

UNIQUENESS OF POSITIVE SOLUTION TO SOME COUPLED COOPERATIVE VARIATIONAL ELLIPTIC SYSTEMS

YULIAN AN, JANN-LONG CHERN, AND JUNPING SHI

ABSTRACT. The uniqueness of positive solutions to some semilinear elliptic systems with variational structure arising from mathematical physics is proved. The key ingredient of the proof is the oscillatory behavior of solutions to linearized equations for cooperative semilinear elliptic systems of two equations on one-dimensional domains, and it is shown that the stability of the positive solutions for such semilinear system is closely related to the oscillatory behavior.

1. INTRODUCTION

Systems of nonlinear elliptic type partial differential equations arise from many models in mathematical physics, such as the nonlinear static Chern-Simons-Higgs equations of classical field theory [9, 12, 16, 19, 20, 21, 59, 60, 68], and standing wave solutions of coupled nonlinear Schrödinger equations from Bose-Einstein condensation [1, 11, 47, 58, 63, 65]. In the case of two interacting particles or waves, the static equation is in form

$$(1.1) \quad \Delta u_1 + f(u_1, u_2) = 0, \quad \Delta u_2 + g(u_1, u_2) = 0, \quad x \in \Omega,$$

where Ω is \mathbb{R}^n or a bounded domain in \mathbb{R}^n . While the existence of positive solutions to (1.1) have been obtained through various variational or other methods, the uniqueness or exact multiplicity of solutions have been mostly open.

In this article, we provide a rather general approach of proving the uniqueness of positive solution to the system in one dimensional space. To achieve that, we prove some general properties of associated linearized system which resembles the classic Sturm comparison principle, and with these properties, for some important systems with a variational structure, we prove the uniqueness of the solution of

$$(1.2) \quad \begin{cases} u_1'' + f(u_1, u_2) = 0, & x \in \mathbb{R}, \\ u_2'' + g(u_1, u_2) = 0, & x \in \mathbb{R}, \\ u_1(x) > 0, \quad u_2(x) > 0, & x \in \mathbb{R}, \\ u_1(x) \rightarrow 0, \quad u_2(x) \rightarrow 0, & |x| \rightarrow \infty, \end{cases}$$

2010 *Mathematics Subject Classification.* Primary 35J47; Secondary 35A02, 35J66, 35J91.
Partially supported by Natural Science Foundation of China (11271261).
Partially supported by MOST of Taiwan under grant no. MOST-104-2115-M-008-010-MY3.
Partially supported by US-NSF grants DMS-1022648 and DMS-1313243.

or the solution of the related Dirichlet boundary value problem:

$$(1.3) \quad \begin{cases} u_1'' + f(u_1, u_2) = 0, & -R < x < R, \\ u_2'' + g(u_1, u_2) = 0, & -R < x < R, \\ u_1(x) > 0, \quad u_2(x) > 0, & -R < x < R, \\ u_1(\pm R) = 0, \quad u_2(\pm R) = 0. \end{cases}$$

Define $\mathbb{R}_+^2 = \{(u_1, u_2) : u_1 \geq 0, u_2 \geq 0\}$. Throughout the paper, we assume that the nonlinear functions f, g in (1.2) and (1.3) satisfy

(f1) $f, g \in C^1(\mathbb{R}_+^2)$;

(f2) (Cooperativeness) Define the Jacobian of the vector field (f, g) to be

$$(1.4) \quad J(u_1, u_2) = \begin{pmatrix} \frac{\partial f}{\partial u_1}(u_1, u_2) & \frac{\partial f}{\partial u_2}(u_1, u_2) \\ \frac{\partial g}{\partial u_1}(u_1, u_2) & \frac{\partial g}{\partial u_2}(u_1, u_2) \end{pmatrix} \equiv \begin{pmatrix} f_1(u_1, u_2) & f_2(u_1, u_2) \\ g_1(u_1, u_2) & g_2(u_1, u_2) \end{pmatrix}.$$

Then (f, g) is said to be *cooperative* if $f_2(u_1, u_2) \geq 0$ and $g_1(u_1, u_2) \geq 0$ for $(u_1, u_2) \in \mathbb{R}_+^2$, and $f_2(u_1, u_2) > 0$ and $g_1(u_1, u_2) > 0$ for $(u_1, u_2) \in \text{int}(\mathbb{R}_+^2)$.

Under the conditions (f1) and (f2), it is well-known that a positive solution $(u_1(x), u_2(x))$ of (1.3) must be an even function in the sense that $u_i(-x) = u_i(x)$ and $u_i'(x) < 0$ for $x \in (0, R)$ (see [64]), and the symmetry properties for positive solutions to (1.2) have also been established in [8, 33, 50] under some additional assumptions on f and g at $(u_1, u_2) = (0, 0)$. These work are natural extensions of the classical results in [37, 38] for the scalar equation since the maximum principle also holds for elliptic systems with cooperative nonlinearities [28, 32, 61].

Our first result is a Sturm comparison type result for positive solutions to system (1.2) or (1.3). We note that the Sturm comparison lemma can be regarded as another aspect of maximum principle in one-dimensional space. A simplified version of the classical Sturm comparison lemma is: suppose that $w_1(x)$ and $w_2(x)$ are two linear independent solutions of $w'' + q(x)w = 0$ where q is continuous on $[a, b]$, and $w_1(a) = w_1(b) = 0$, then w_2 has a zero in (a, b) . A straightforward application of this lemma is for a solution of

$$(1.5) \quad u'' + g(u) = 0, \quad x \in (0, R), \quad u'(0) = 0, \quad u(0) = \alpha,$$

such that $u(x) > 0$, $u'(x) < 0$ in $(0, R)$, then any solution ϕ of the linearized equation

$$(1.6) \quad \phi'' + g'(u(x))\phi = 0, \quad x \in (0, R)$$

changes sign at most once in $(0, R)$ since $u'(x)$ is also a solution of (1.6), $u'(0) = 0$ and $u'(x) < 0$ in $(0, R)$. Our result for solutions of linearized equation around a positive solution to (1.2) or (1.3) resembles the one above for the scalar equation. More precisely, suppose that f, g satisfy (f1) and (f2), (u_1, u_2) is a positive solution of (1.2) or (1.3), and f_1, f_2, g_1, g_2 are as defined in (1.4), we prove that if (ϕ_1, ψ_1) (resp. (ϕ_2, ψ_2)) is a solution of

$$(1.7) \quad \begin{cases} \phi'' + f_1\phi + f_2\psi = 0, & 0 < x < R, \\ \psi'' + g_1\phi + g_2\psi = 0, & 0 < x < R, \\ \phi'(0) = \psi'(0) = 0, \\ (\phi(0), \psi(0)) = (1, 0), \quad (\text{resp. } (0, 1)), \end{cases}$$

then each of ϕ_i and ψ_i ($i = 1, 2$) changes sign at most once in $(0, R)$ (see Lemma 2.2 and Corollary 2.3 for more precise statements). It is well-known that oscillatory properties of solutions to the linearized equation is critical for the stability and uniqueness of positive solution of semilinear elliptic equations [3, 44, 53, 54], hence the non-oscillatory property for solutions of (1.7) is very useful for the stability and uniqueness of positive solutions to (1.2) or (1.3). We remark that such property usually does not hold for higher dimensional radial Laplacian $Lu = r^{1-n}(r^{n-1}u)'$ even for the scalar case, hence the spatial dimension $n = 1$ is a critical assumption here.

Our second result is related to the stability of a positive solution to (1.2) or (1.3). It is known that for (1.5), the number of sign-changes of the solution of linearized equation is related to stability of a positive solution. If the solution of linearized equation does not change sign, then the positive solution is linearly stable; while the solution of linearized equation changes sign once, then the positive solution is linearly unstable. Here we also establish such a connection between the number of sign-changes of the solution to the linearized equation (1.7) and the stability of a positive solution of the system (1.2). More precisely we show that, suppose that f, g satisfy (f1) and (f2), a positive solution (u_1, u_2) of (1.3) is stable if there exists a real number $c > 0$ such that $\phi_1 + c\phi_2 > 0$ and $\psi_1 + c\psi_2 > 0$ in $(0, R]$, where (ϕ_i, ψ_i) is the solution of (1.7); and (u_1, u_2) is unstable if and only if for all $c \geq 0$, at least one of $\phi_1 + c\phi_2$ or $\psi_1 + c\psi_2$ is not positive in $(0, R)$ (see Proposition 2.8).

The non-oscillatory results above are proved under rather general conditions (f1) and (f2) on (f, g) , and these results pave the way for the stability, non-degeneracy and uniqueness of the positive solution to (1.2) or (1.3) from a wide range of applications. Two additional structures on (f, g) would be needed for these further results: (i) the growth rate of functions f and g ; and (ii) a variational structure for the vector field (f, g) .

The growth rate of the functions f and g plays an important role in the qualitative behavior of the solutions to (1.2) and (1.3). Here we define several conditions on the growth rate of f and g :

(f3) (Superlinear) The vector field (f, g) is said to be *superlinear* if for all $(u_1, u_2) \in \mathbb{R}_+^2$,

$$(1.8) \quad f_1 u_1 + f_2 u_2 - f \geq 0, \quad g_1 u_1 + g_2 u_2 - g \geq 0;$$

(f3') (Strongly superlinear) The vector field (f, g) is said to be *strongly superlinear*, if for all $(u_1, u_2) \in \mathbb{R}_+^2$,

$$(1.9) \quad f_1 u_1 - f \geq 0, \quad g_2 u_2 - g \geq 0;$$

(f4) (Sublinear) The vector field (f, g) is said to be *sublinear* if for all $(u_1, u_2) \in \mathbb{R}_+^2$,

$$(1.10) \quad f_1 u_1 + f_2 u_2 - f \leq 0, \quad g_1 u_1 + g_2 u_2 - g \leq 0;$$

(f4') (Weakly sublinear) The vector field (f, g) is said to be *weakly sublinear*, if for all $(u_1, u_2) \in \mathbb{R}_+^2$,

$$(1.11) \quad f_1 u_1 - f \leq 0, \quad g_2 u_2 - g \leq 0.$$

We remark that all definitions above are actually for a single function $f : \mathbb{R}_+^2 \rightarrow \mathbb{R}$, but we assume that f and g have the same type of growth rate in this article. Note that under the cooperativeness assumption (f2), condition (f3') implies (f3), while

condition (f4) implies (f4'), which is the reason for the “strongly” and “weakly” in the definition. On the other hand, we notice that a function f can be both weakly sublinear and superlinear. A prototypical example of a cooperative type function f is

$$(1.12) \quad f(u_1, u_2) = \lambda u_1 + \mu u_2 + \sum_{i=1}^k a_i u_1^{p_i} u_2^{q_i},$$

where $\lambda \in \mathbb{R}$, $\mu \in [0, \infty)$, $a_i > 0$, and $p_i, q_i \geq 0$ for all $1 \leq i \leq k$. Under these general conditions, it is easy to verify that

$p_i + q_i \geq 1 \Rightarrow f$ is superlinear; $p_i \geq 1$, $\mu = 0 \Rightarrow f$ is strongly superlinear;

$p_i + q_i \leq 1 \Rightarrow f$ is sublinear; $p_i \leq 1 \Rightarrow f$ is weakly sublinear.

Hence f is both weakly sublinear and superlinear if $0 \leq p_i \leq 1$, $0 \leq q_i$ and $p_i + q_i \geq 1$, which always holds when $p_i = 1$. The notion of superlinear and sublinear growth rate for a uni-variable function $g : \mathbb{R}_+ \rightarrow \mathbb{R}$ was considered in [54], and definition here can be considered as a generalization of the definition in [54] to multi-variable functions. Similar definitions of multi-variable sublinear/superlinear functions can also be found in [18, 25, 57]. It is known that sublinear/superlinear properties are related to the stability of positive solutions of (1.3). We prove that when (f, g) is sublinear, then any positive solution of (1.3) is stable, while when (f, g) is superlinear, then any positive solution of (1.3) is unstable (see Lemma 2.5 and also [25]).

The final assumption for non-degeneracy and uniqueness of positive solution is the variational structure on the vector field (f, g) . Here two possible variational structure can be defined as in [29, 30]. The system (1.2) or (1.3) is a *Hamiltonian system* if there exists a differentiable function $H(u_1, u_2)$ such that

$$(1.13) \quad f(u_1, u_2) = \frac{\partial H(u_1, u_2)}{\partial u_2}, \quad \text{and} \quad g(u_1, u_2) = \frac{\partial H(u_1, u_2)}{\partial u_1};$$

Clearly a Hamiltonian system satisfies $f_1 = g_2$. For a Hamiltonian system, if $(u_1(x), u_2(x))$ is a solution of (2.3), we define

$$(1.14) \quad H_0(x) = u_1'(x)u_2'(x) + H(u_1(x), u_2(x)),$$

then $H_0'(x) = 0$ hence $H_0(x) \equiv H_0(0)$ for $x > 0$. On the other hand, the system (1.2) or (1.3) is a *gradient system* if there exists a differentiable function $F(u_1, u_2)$ such that

$$(1.15) \quad f(u_1, u_2) = \frac{\partial F(u_1, u_2)}{\partial u_1}, \quad \text{and} \quad g(u_1, u_2) = \frac{\partial F(u_1, u_2)}{\partial u_2}.$$

Clearly a gradient system satisfies $f_2 = g_1$ hence the Jacobian matrix is symmetric and the corresponding linearized equation is self-adjoint. For a gradient system, if $(u_1(x), u_2(x))$ is a solution of (2.3), we define

$$(1.16) \quad F_0(x) = \frac{1}{2}[u_1'(x)]^2 + \frac{1}{2}[u_2'(x)]^2 + F(u_1(x), u_2(x)),$$

then $F_0'(x) = 0$ hence $F_0(x) \equiv F_0(0)$ for $x > 0$. Both the energy functions H_0 and F_0 are generalizations of the energy function $G_0(x) = \frac{1}{2}[u'(x)]^2 + G(u(x))$ for (1.5) where $G(u) = \int_0^u g(s)ds$. For the scalar equation

$$(1.17) \quad u'' + g(u) = 0, \quad x \in \mathbb{R}, \quad u'(0) = 0, \quad \lim_{|x| \rightarrow \infty} u(x) = 0,$$

the energy function G_0 alone can guarantee the uniqueness of positive solution to (1.17), which in general is not the case for the system (1.3). But with the Hamiltonian of Gradient structure, the uniqueness of positive solution to (1.3) can be proved by combining with oscillatory property.

Our third key result is that assume (u_1, u_2) is a positive solution of (1.3), (f, g) satisfies (f1) and (f2), (f, g) is superlinear (thus (u_1, u_2) is unstable), and in addition, if (f, g) is weakly sublinear, and it is a Hamiltonian or gradient system, then (u_1, u_2) must be non-degenerate, which often suggests uniqueness. Combining the cooperative and variational structure, the weakly sublinear and superlinear properties, we can prove that the positive solution to (1.3) for certain (f, g) is unique for any given $R > 0$, and it also implies the uniqueness of positive solution to (1.2) when it exists. More precisely it can be shown that all solutions of (1.3) can be represented by a curve $N = \{(\alpha, \beta(\alpha), R(\alpha)) : \alpha \in A\}$ where A is a subset of \mathbb{R}_+ . For a given α in the admissible set A , there is a unique $\beta(\alpha) > 0$ such that (1.3) has a unique positive solution (u_1, u_2) satisfying $u_1(0) = \alpha$, $u_2(0) = \beta(\alpha)$ and $R = R(\alpha)$. Moreover it shown that $\beta(\alpha)$ is a strictly increasing function of α , and $R(\alpha)$ is a strictly increasing (decreasing) function of α when (f, g) is sublinear (superlinear). The monotonicity of $R(\alpha)$ implies the uniqueness of positive solution to (1.3) for a given $R > 0$ (see Section 3 for more precise results).

In Section 3, this program of proving uniqueness of positive solution to (1.3) is achieved for

- (A) (Schrödinger type [31, 33, 39]) $f(u_1, u_2) = -u_1 + u_2^q$, $g(u_1, u_2) = -u_2 + u_1^p$, where $p, q > 1$.
- (B) (optics model [48, 49, 69]) $f(u_1, u_2) = -bu_1 + u_1u_2$, $g(u_1, u_2) = -cu_2 + u_1^2/2$, where $b, c > 0$.

Here system (A) is a Hamiltonian system, and the one in (B) is a gradient system. The more general results established in Section 2 can be applied to prove the uniqueness of positive solution of (1.3) as long as (f, g) is (i) cooperative, (ii) weakly sublinear and superlinear, and (iii) Hamiltonian or gradient.

The uniqueness of positive solution of (1.3) for sublinear (f, g) case holds for a general bounded domain in \mathbb{R}^n with $n \geq 1$, see [4, 18, 25]. Some methods for the uniqueness of the positive solution of semilinear elliptic systems used in this paper were also used in some of our earlier work [13, 15, 16, 17, 19, 20], but the oscillatory results for the linearized equation here are the most general ones as they do not rely on specific algebraic form of the nonlinearities f and g . We expect these results will be useful for the uniqueness of the positive solution to other cooperative systems. Previously the uniqueness of the positive solution was mostly only known for the Lane-Emden system in which the nonlinearities are power functions, see [26, 27, 41, 43, 56]. Recently the uniqueness of positive solution to the coupled Schrödinger equations in some special cases was proved in [11, 40, 51, 67].

The remaining part of the paper is organized as follows. In Section 2, we consider the properties of solutions to linearized equations, and the relation between the stability of solution and nodal properties of solutions to linearized equations is considered. The uniqueness of positive solution to several semilinear cooperative elliptic systems with weakly sublinear and superlinear nonlinearities are studied in Section 3.

2. LINEARIZED EQUATIONS AND STABILITY

2.1. Preliminaries. In the following we always assume that (f, g) satisfies the condition (f1) and (f2). With a possible translation, a solution of (1.2) or (1.3) is necessarily symmetric with respect to $x = 0$ and decreasing when $x > 0$ [8, 64], hence it satisfies either

$$(2.1) \quad \begin{cases} u_1'' + f(u_1, u_2) = 0, & 0 < x < \infty, \\ u_2'' + g(u_1, u_2) = 0, & 0 < x < \infty, \\ u_1(x) > 0, \quad u_2(x) > 0, \quad u_1'(x) < 0, \quad u_2'(x) < 0, & 0 < x < \infty, \\ u_1'(0) = u_2'(0) = 0, \\ u_1(x) \rightarrow 0, \quad u_2(x) \rightarrow 0, & x \rightarrow \infty, \end{cases}$$

or

$$(2.2) \quad \begin{cases} u_1'' + f(u_1, u_2) = 0, & 0 < x < R, \\ u_2'' + g(u_1, u_2) = 0, & 0 < x < R, \\ u_1(x) > 0, \quad u_2(x) > 0, \quad u_1'(x) < 0, \quad u_2'(x) < 0, & 0 < x < R, \\ u_1'(0) = u_2'(0) = 0, \\ u_1(R) = 0, \quad u_2(R) = 0. \end{cases}$$

A solution $(u_1(x), u_2(x))$ of (2.1) or (2.2) is a solution of the initial value problem:

$$(2.3) \quad \begin{cases} u_1'' + f(u_1, u_2) = 0, & x > 0, \\ u_2'' + g(u_1, u_2) = 0, & x > 0, \\ u_1'(0) = u_2'(0) = 0, \\ u_1(0) = \alpha, \quad u_2(0) = \beta, \end{cases}$$

where $\alpha > 0$ and $\beta > 0$. The local existence and uniqueness of the solution of (2.3) can be proved via a standard argument. We denote the solution of (2.3) by $(u_1(x; \alpha, \beta), u_2(x; \alpha, \beta))$ or simply $(u_1(x), u_2(x))$ when there is no confusion. We will only consider a solution of (2.3) satisfying $u_i(x) > 0$ and $u_i'(x) < 0$ for $x \in (0, \delta)$, $i = 1, 2$. This requires the initial value (α, β) to satisfy $f(\alpha, \beta) \geq 0$ and $g(\alpha, \beta) \geq 0$, and at least one of them is strictly greater than zero. A solution of (2.2) is a *crossing solution* of (2.3), and a solution of (2.1) is a *ground state solution*. The local solution $(u_1(x), u_2(x))$ of (2.3) can be extended to a maximal interval $(0, R(\alpha, \beta))$ so that $u_i(x) > 0$ and $u_i'(x) < 0$ in $(0, R(\alpha, \beta))$, $i = 1, 2$. In the following we will use $R = R(\alpha, \beta)$ when there is no confusion.

If $R < \infty$, then, for technical reason, we need to extend the solution beyond $x = R$. We extend the definition of (f, g) to \mathbb{R}^2 so that $f, g \in C^1(\mathbb{R}^2)$. Hence the solution $(u_1(x), u_2(x))$ can be extended to $x \in [0, R + \varepsilon]$ for a positive $\varepsilon = \varepsilon(\alpha, \beta)$ with u_i or $-u_i'$ possibly negative in $(R, R + \varepsilon]$. In the following we shall always assume that the domain of a solution $(u_1(x), u_2(x))$ is $x \in [0, R + \varepsilon]$ when $R < \infty$.

To consider the dependence of solutions on the initial values, we consider the linearized equation of $(u_1(x), u_2(x))$ with respect to the initial value (α, β) . Let $W(x)$ be a 2×2 matrix function defined as

$$(2.4) \quad W(x) \equiv \begin{pmatrix} \frac{\partial u_1(x; \alpha, \beta)}{\partial \alpha} & \frac{\partial u_1(x; \alpha, \beta)}{\partial \beta} \\ \frac{\partial u_2(x; \alpha, \beta)}{\partial \alpha} & \frac{\partial u_2(x; \alpha, \beta)}{\partial \beta} \end{pmatrix}.$$

Then $W(x)$ satisfies

$$(2.5) \quad \begin{cases} W'' + J(u_1(x), u_2(x))W = 0, & x > 0, \\ W(0) = I, \quad W'(0) = 0. \end{cases}$$

Here I is the identity matrix, J is defined in (1.4), W' and W'' are the matrices of derivatives of each entry W_{ij} of W with respect to x . Equivalently we can write the componentwise equation: let $(\phi_1, \psi_1) = \left(\frac{\partial u_1(x; \alpha, \beta)}{\partial \alpha}, \frac{\partial u_2(x; \alpha, \beta)}{\partial \alpha} \right)$, then (ϕ_1, ψ_1) satisfies

$$(2.6) \quad \begin{cases} \phi_1'' + f_1\phi_1 + f_2\psi_1 = 0, & x > 0, \\ \psi_1'' + g_1\phi_1 + g_2\psi_1 = 0, & x > 0, \\ \phi_1(0) = 1, \quad \phi_1'(0) = 0, \\ \psi_1(0) = 0, \quad \psi_1'(0) = 0; \end{cases}$$

and let $(\phi_2, \psi_2) = \left(\frac{\partial u_1(x; \alpha, \beta)}{\partial \beta}, \frac{\partial u_2(x; \alpha, \beta)}{\partial \beta} \right)$, then (ϕ_2, ψ_2) satisfies

$$(2.7) \quad \begin{cases} \phi_2'' + f_1\phi_2 + f_2\psi_2 = 0, & x > 0, \\ \psi_2'' + g_1\phi_2 + g_2\psi_2 = 0, & x > 0, \\ \phi_2(0) = 0, \quad \phi_2'(0) = 0, \\ \psi_2(0) = 1, \quad \psi_2'(0) = 0. \end{cases}$$

To study the oscillatory behavior of (ϕ_i, ψ_i) ($i = 1, 2$), we first consider two auxiliary equations. Let (ϕ_3, ψ_3) be the unique solution of

$$(2.8) \quad \begin{cases} \phi_3'' + f_1\phi_3 = 0, & x > 0, \\ \psi_3'' + g_2\psi_3 = 0, & x > 0, \\ \phi_3(0) = 1, \quad \phi_3'(0) = 0, \\ \psi_3(0) = 1, \quad \psi_3'(0) = 0. \end{cases}$$

The oscillatory behavior of (ϕ_3, ψ_3) can be obtained as follows.

Lemma 2.1. *Let (u_1, u_2) be a solution of (2.3) such that $u_1(x) > 0$, $u_2(x) > 0$, $u_1'(x) < 0$ and $u_2'(x) < 0$ for $x \in (0, R)$ (R may be ∞), and let (ϕ_3, ψ_3) be defined as in (2.8). Assume that (f, g) is cooperative as defined in (f2). Then*

- (1) *Each of ϕ_3 and ψ_3 has at most one zero in $(0, R]$;*
- (2) *If in addition, (f, g) is weakly sublinear, and (u_1, u_2) is a solution of (2.2), then each of ϕ_3 and ψ_3 is positive in $(0, R]$;*
- (3) *If in addition, (f, g) is strongly superlinear, and (u_1, u_2) is a solution of (2.2), then each of ϕ_3 and ψ_3 changes sign exactly once in $(0, R)$, and $\phi_3(R) < 0$, $\psi_3(R) < 0$.*

Proof. (1) We only prove it for ϕ_3 , and the proof for ψ_3 is similar. Suppose that ϕ_3 changes sign more than once. Let $0 < x_1 < x_2 \leq R$ be the first two zeros of ϕ_3 . From the equation which ϕ_3 satisfies, each zero of ϕ_3 is a simple one. Then $\phi_3(x) < 0$ in (x_1, x_2) , $\phi_3'(x_1) < 0$ and $\phi_3'(x_2) > 0$. Notice that (u_1', u_2') satisfies

$$(2.9) \quad \begin{cases} (u_1')'' + f_1u_1' + f_2u_2' = 0, & x > 0, \\ (u_2')'' + g_1u_1' + g_2u_2' = 0, & x > 0, \\ u_1'(0) = 0, \quad u_2'(0) = 0. \end{cases}$$

Then multiplying the equation of ϕ_3 in (2.8) by u_1' , multiplying the equation of u_1' in (2.9) by ϕ_3 , subtracting and integrating on (x_1, x_2) , we obtain

$$(2.10) \quad \phi_3'(x_2)u_1'(x_2) - \phi_3'(x_1)u_1'(x_1) = \int_{x_1}^{x_2} f_2\phi_3u_2'dx.$$

The left hand side of (2.10) is non-positive since $u_1'(x) < 0$ in $(0, R)$, while the right hand side of (2.10) is positive since $f_2 > 0$, $\phi_3 < 0$ and $u_2' < 0$ in (x_1, x_2) . That is a contradiction. Hence ϕ_3 cannot have more than one zeros in $(0, R]$.

(2) Next we assume that (f, g) is weakly sublinear, and (u_1, u_2) is a solution of (2.2). Suppose that ϕ_3 has a zero $x_1 \in (0, R]$, then $\phi_3(x) > 0$ for $x \in (0, x_1)$ and $\phi_3(x_1) = 0$. We rewrite the equations in (2.2) to

$$(2.11) \quad \begin{cases} u_1'' + f_1u_1 = f_1u_1 - f, & x > 0, \\ u_2'' + g_2u_2 = g_2u_2 - g, & x > 0. \end{cases}$$

Multiplying the equation of ϕ_3 in (2.8) by u_1 , multiplying the equation of u_1 in (2.11) by ϕ_3 , subtracting and integrating on $(0, x_1)$, we have

$$(2.12) \quad \phi_3'(x_1)u_1(x_1) = - \int_0^{x_1} (f_1u_1 - f)\phi_3dx.$$

Then the left hand side of (2.12) is negative, while the right hand side of (2.12) is non-negative. This is a contradiction, so $\phi_3(x) > 0$ for $x \in (0, R]$.

(3) Now we also assume (f, g) is strongly superlinear, and (u_1, u_2) is a solution of (2.2). Suppose that ϕ_3 does not change sign, then $\phi_3(x) > 0$ for $x \in (0, R)$. Similar to (2), we have

$$(2.13) \quad -\phi_3(R)u_1'(R) = - \int_0^R (f_1u_1 - f)\phi_3dx.$$

Then the left hand side of (2.13) is non-negative, while the right hand side of (2.13) is negative. This contradiction implies that ϕ_3 has to change sign in $(0, R)$, and $\phi_3(R) < 0$ since ϕ_3 cannot have more than one zeros from the previous proof. \square

The following result provides a key oscillatory result for the solutions of the linearized equation (2.6) and (2.7).

Lemma 2.2. *Let (u_1, u_2) be a solution of (2.3) such that $u_1(x) > 0$, $u_2(x) > 0$ for $x \in (0, R)$, and $u_1'(x) < 0$ and $u_2'(x) < 0$ for $x \in (0, R]$ (when R is ∞ , then $u_i'(x) < 0$ for $x \in (0, R)$), and let ϕ_i, ψ_i ($i = 1, 2, 3$) be defined as in (2.6), (2.7) and (2.8). Assume that (f, g) is cooperative as defined in (f2). Then*

- (1) $\phi_1(x)$ changes sign at most once, and $\psi_1(x) < 0$ for $x \in (0, R)$; if in addition $\phi_3(x) > 0$, then $\phi_1(x) > 0$ for $x \in (0, R)$;
- (2) $\psi_2(x)$ changes sign at most once, and $\phi_2(x) < 0$ for $x \in (0, R)$; if in addition $\psi_3(x) > 0$, then $\psi_2(x) > 0$ for $x \in (0, R)$.

Proof. If $R < \infty$, then from earlier comment, we extend the definition of the solution to $[0, R + \varepsilon]$ and since $u_i'(R) < 0$, then we can assume that $u_i'(x) < 0$ for $x \in (0, R + \varepsilon]$. In the following we only prove the result for ϕ_1 and ψ_1 , and the proof for the other case is similar. Since $\phi_1(0) > 0$, $\psi_1(0) = 0$, $\psi_1'(0) = 0$, and $\psi_1''(0) = -g_2\psi_1(0) - g_1\phi_1(0) < 0$. Then for some $x_0 > 0$, $\phi_1(x) > 0$ and $\psi_1(x) < 0$ in $(0, x_0)$. Define

$$(2.14) \quad x_1 = \sup\{0 < x < R : \phi_1(r) > 0 \text{ and } \psi_1(r) < 0 \text{ in } (0, x)\}.$$

If $x_1 = R$, then the result holds. So we assume $x_1 < R$. Then either $\phi_1(x_1) = 0$ or $\psi_1(x_1) = 0$. If $\psi_1(x_1) = 0$, then $\psi_1(x) < 0$ in $(0, x_1)$ and $\psi_1'(x_1) \geq 0$. Then multiplying the equation of ψ_1 in (2.6) by u_2' , multiplying the equation of u_2' in (2.9) by ψ_1 , subtracting and integrating on $(0, x_1)$, we obtain

$$(2.15) \quad [\psi_1' u_2' - \psi_1 u_2'']|_0^{x_1} = - \int_0^{x_1} g_1(\phi_1 u_2' - \psi_1 u_1') dx.$$

The left hand side is $\psi_1'(x_1)u_2'(x_1) \leq 0$, and the right hand side is positive since $g_1 > 0$, $\phi_1 > 0$, $\psi_1 < 0$, $u_1' < 0$ and $u_2' < 0$ in $(0, x_1)$. That is a contradiction.

Hence $\psi_1(x_1) = 0$ does not hold and one must have $\phi_1(x_1) = 0$, which implies that $\psi_1(x) < 0$ in $(0, x_1]$, $\phi_1(x) > 0$ in $(0, x_1)$ and $\phi_1'(x_1) \leq 0$. We claim that in this case ϕ_3 defined in (2.8) changes sign in $(0, x_1)$. If not, $\phi_3(x) > 0$ in $(0, x_1)$, then from the equation of ϕ_1 and ϕ_3 , we obtain that

$$(2.16) \quad \phi_3(x_1)\phi_1'(x_1) = - \int_0^{x_1} f_2\psi_1\phi_3 dx,$$

which is contradiction since the left hand side is non-positive, and the right hand side is positive. Hence ϕ_3 has to change sign before ϕ_1 does. From Lemma 2.1, ϕ_3 changes sign at most once in $(0, R)$. If ϕ_3 does not change sign in $(0, R)$, then ϕ_1 and ψ_1 do not change sign either, thus the conclusion in the lemma holds.

So now we assume that ϕ_3 changes sign exactly once at $x_0 \in (0, R)$, then $0 < x_0 < x_1$, and $\phi_3(x) < 0$ in $[x_0, R]$. We claim that ϕ_1 cannot have more than one zeros in $(0, R)$. We define another pair of auxiliary functions $(\phi_{4,k}, \psi_{4,k})$ which is defined by $\phi_{4,k} = \phi_1 + ku_1'$ and $\psi_{4,k} = \psi_1 + ku_2'$ for $k \geq 0$. (ϕ_1, ψ_1) and (u_1', u_2') satisfy the same linear equations, so does $(\phi_{4,k}, \psi_{4,k})$, namely

$$(2.17) \quad \begin{cases} \phi_{4,k}'' + f_1\phi_{4,k} + f_2\psi_{4,k} = 0, & x > 0, \\ \psi_{4,k}'' + g_1\phi_{4,k} + g_2\psi_{4,k} = 0, & x > 0, \\ \phi_{4,k}(0) = 1, \quad \psi_{4,k}(0) = 0. \end{cases}$$

When $k > 0$ is large enough, $\psi_{4,k}(x) < 0$ for $x \in (0, R)$, and $\phi_{4,k}$ has exactly one zero $x_{1,k} \in (0, R)$ which approaches to 0 as $k \rightarrow \infty$ since $\phi_{4,k}'(0) = ku_1''(0) < 0$. Hence we can define

$$(2.18) \quad k_* = \inf\{k_0 > 0 : \psi_{4,k}(x) < 0 \text{ for } x \in (0, R), \\ \text{and } \phi_{4,k} \text{ has exactly one zero } x_{1,k} \in (0, R) \text{ for all } k \geq k_0\}.$$

Such k_* clearly exists as for any $k < 0$, $\psi_{4,k}(x) > 0$ near $x = 0$. Hence $k_* \geq 0$. Suppose that $k_* > 0$. Then $\phi_{4,k_*}(x) > 0$ and $\psi_{4,k_*}(x) < 0$ for small x , and one of the following alternatives occurs:

- (i) there exists $x_2 > 0$ such that $\psi_{4,k_*}(x) < 0$ in $(0, x_2)$ and $\psi_{4,k_*}(x_2) = 0$;
- (ii) ϕ_{4,k_*} has more than one zeros in $(0, R)$;
- (iii) $\phi_{4,k}$ has exactly one zero $x_{1,k} \in (0, R)$ for $k > k_*$, but $\phi_{4,k}$ has no zero in $(0, R)$ when $k = k_*$.

If case (iii) occurs, then $\phi_1(x) > 0$ for $x \in (0, R)$ since $\phi_{4,k}$ decreases in k , then the conclusion in the lemma holds. It is obvious that if (iii) occurs, then (ii) cannot occur. Also notice that if (iii) occurs, then (i) cannot occur. On the contrary, suppose case (i) and case (iii) both occur. Similar to (2.15), if a, b are two consecutive

zeros of $\psi_{4,k}$, then

$$(2.19) \quad [\psi'_{4,k}u'_2 - \psi_{4,k}u''_2] \Big|_a^b = - \int_a^b g_1(\phi_{4,k}u'_2 - \psi_{4,k}u'_1)dx.$$

For $k = k_*$, let $a = 0$ and $b = x_2$. Then the right side of (2.19) is positive and the left side is nonnegative. This is a contradiction.

Suppose that case (i) occurs but (ii) does not. Similar to the discussion of last paragraph, we must have $x_2 > x_{1,k_*}$, the unique zero of ϕ_{4,k_*} . In particular this implies that $\phi_{4,k_*}(x_2) < 0$. Also x_2 is necessarily a local maximum point of ψ_{4,k_*} . However, from (2.17),

$$\psi''_{4,k_*}(x_2) = -g_1\phi_{4,k_*}(x_2) > 0.$$

we reach another contradiction.

Next we consider the case (ii) occurs but not (i), hence $\psi_{4,k_*}(x) < 0$ in $(0, R)$. Let $x_3 < x_4$ be the first two zeros of ϕ_{4,k_*} . By using the equations of $\phi_{4,k}$ and u'_1 , if a, b are two consecutive zeros of $\phi_{4,k}$, then

$$(2.20) \quad [\phi'_{4,k}u'_1 - \phi_{4,k}u''_1] \Big|_a^b = - \int_a^b f_2(\psi_{4,k}u'_1 - \phi_{4,k}u'_2)dx.$$

Since $\phi_{4,k_*}(0) = 1 > 0$, then $\phi_{4,k_*}(x) < 0$ in (x_3, x_4) , and x_4 is necessarily a local maximum point. However, from (2.17),

$$\phi''_{4,k_*}(x_4) = -f_2\psi_{4,k_*}(x_4) > 0.$$

Contradiction.

Finally we consider the case (i) and (ii) occur simultaneously, that is ϕ_{4,k_*} has the first two zeros $x_5 < x_6$, and ψ_{4,k_*} has a first zero at x_2 . Again we have $\phi_{4,k_*}(x) < 0$ in (x_5, x_6) . Apparently $x_2 > x_5$ from proof above. If $x_5 < x_2 < x_6$, then the argument for case (i) still works, and we can use (2.19) to get a contradiction; if $x_2 > x_6$, then the argument for case (ii) still works, and we can use (2.20) to get contradiction. Finally if $x_2 = x_6$, we will have $\phi_{4,k_*}(x_2) = \phi'_{4,k_*}(x_2) = 0$ and $\psi_{4,k_*}(x_2) = \psi'_{4,k_*}(x_2) = 0$, but that would implies $\phi_{4,k_*} \equiv \psi_{4,k_*} \equiv 0$ for $x \in (0, R)$ from the uniqueness of second order linear ODE.

Since we reach contradictions in all the cases (i) and (ii), then we cannot have $k_* > 0$. Therefore $k_* = 0$ and the proof is completed. \square

For the weakly sublinear case, Lemmas 2.1 and 2.2 imply the following result which will be used later.

Corollary 2.3. Let (u_1, u_2) be a solution of (2.2) such that $u'_1(R) < 0$ and $u'_2(R) < 0$, and let (ϕ_i, ψ_i) ($i = 1, 2, 3$) be the solution of (2.6), (2.7) and (2.8) respectively. Assume that (f, g) is cooperative as defined in (f2) and is weakly sublinear, then both of ϕ_1 and ψ_2 are positive on $[0, R]$.

2.2. Stability. Let (u_1, u_2) be a solution of (1.3). The stability of (u_1, u_2) is determined by the eigenvalue problem:

$$(2.21) \quad \begin{cases} \xi'' + f_1\xi + f_2\eta = -\mu\xi, & x \in I, \\ \eta'' + g_1\xi + g_2\eta = -\mu\eta, & x \in I, \\ \xi(x) = 0, \quad \eta(x) = 0, & x \in \partial I. \end{cases}$$

Here $I = (-R, R)$. Notice that (2.21) can be written as

$$(2.22) \quad L\mathbf{u} = H\mathbf{u} + \mu\mathbf{u},$$

where

$$(2.23) \quad \mathbf{u} = \begin{pmatrix} \xi \\ \eta \end{pmatrix}, \quad L\mathbf{u} = \begin{pmatrix} -\xi'' - f_1\xi \\ -\eta'' - g_2\eta \end{pmatrix}, \quad \text{and } H = \begin{pmatrix} 0 & f_2 \\ g_1 & 0 \end{pmatrix}.$$

Again we assume that (f, g) is cooperative, then the system (2.22) and (2.23) is a linear elliptic system of cooperative type, and the maximum principles hold for such systems. Here we recall some known results:

Lemma 2.4. *Suppose that L, H are given in (2.23), $\mathbf{u} \in X \equiv [W^{2,2}(I) \cap W_0^{1,2}(I)]^2$, and (f, g) is cooperative as defined in (f2). Then*

- (1) $\mu_1 = \inf\{\mu \in \text{Re}(\text{spt}(L - H))\}$ is a real-value eigenvalue of $L - H$, where $\text{spt}(L - H)$ is the spectrum of $L - H$.
- (2) For $\mu = \mu_1$, there exists a unique (up a constant multiple) eigenfunction $\mathbf{u}_1 \in Y \equiv [L^2(I)]^2$, and $\mathbf{u}_1 > 0$ in I .
- (3) For $\mu < \mu_1$, the equation $L\mathbf{u} = H\mathbf{u} + \mu\mathbf{u} + \mathbf{f}$ is uniquely solvable for any $\mathbf{f} \in Y$, and $\mathbf{u} > 0$ as long as $\mathbf{f} \geq 0$.
- (4) (Maximum principle) For $\mu \leq \mu_1$, suppose that $\mathbf{u} \in [W^{2,2}(I)]^2$, satisfies $L\mathbf{u} \geq H\mathbf{u} + \mu\mathbf{u}$ in I , $\mathbf{u} \geq 0$ on ∂I , then $\mathbf{u} \geq 0$ in I .
- (5) If there exists $\mathbf{u} \in [W^{2,2}(I)]^2$, satisfying $L\mathbf{u} \geq H\mathbf{u}$ and $\mathbf{u} \geq 0$ in I , and either $\mathbf{u} \not\equiv 0$ on ∂I or $L\mathbf{u} \not\equiv H\mathbf{u}$ in I , then $\mu_1 > 0$.
- (6) The principal eigenvalue can be characterized by

$$(2.24) \quad \mu_1 = \sup\{\mu : \exists \mathbf{w} \in [W_{loc}^{2,1}(I)]^2 \text{ such that } (L - H)\mathbf{w} \geq \mu\mathbf{w}, \text{ and } \mathbf{w} > 0\}.$$

For the result and proofs parts 1-5 in Lemma 2.4, see Sweers [61] Proposition 3.1 and Theorems 1.1, and the one in part 6, see Birindelli et.al. [7] Theorem 1 (which extends the result in Berestycki et.al. [6] for scalar equations). Moreover from a standard compactness argument, the linear operator $L - H$ has countably many eigenvalues μ_i ($i = 1, 2, \dots$), and the eigenvalues can be ordered so that $\text{Re}(\mu_i - \mu_1) \rightarrow \infty$ as $i \rightarrow \infty$. A solution (u_1, u_2) of (1.3) is called *stable* if $\mu_1 > 0$, *neutrally stable* if $\mu_1 = 0$, and it is *unstable* if $\mu_1 < 0$. The *Morse index* $M(u_1, u_2)$ of (u_1, u_2) is the number of eigenvalues with negative real parts counting multiplicity. Also (u_1, u_2) is *nondegenerate* if $\mu_i \neq 0$ for all $i = 1, 2, \dots$.

One useful fact is that, from part 6 of Lemma 2.4, if (f, g) is cooperative as defined in (f2), then (2.24) implies that for the same (f_1, f_2, g_1, g_2) defined for $I = (-R_1, R_1)$,

$$(2.25) \quad \mu_1(R_1) \leq \mu_1(R_2) \text{ if } R_1 > R_2 > 0,$$

where $\mu_1(R_j)$ is the corresponding principal eigenvalue of (2.21) with $I = (-R, R)$ and $R = R_j$ ($j = 1, 2$).

For a given solution (u_1, u_2) of (1.3), we can also define another eigenvalue problem:

$$(2.26) \quad \begin{cases} \xi'' + f_1\xi + f_2\eta = -\mu\xi, & x \in (0, R), \\ \eta'' + g_1\xi + g_2\eta = -\mu\eta, & x \in (0, R), \\ \xi'(0) = \xi(R) = 0, \quad \eta'(0) = \eta(R) = 0. \end{cases}$$

If (μ, ξ, η) is an eigen-triple of (2.26), then it also satisfies (2.21) via an even extension from $[0, R]$ to $[-R, R]$. This also implies that if (2.26) has a positive eigenvector (ξ_1, η_1) for an eigenvalue μ_1 , then extended (μ_1, ξ_1, η_1) is also the principal eigen-triple for (2.21). On the other hand, if (μ_1, ξ_1, η_1) is the principal eigen-triple for (2.21) such that $\xi_1, \eta_1 > 0$, then $(\xi_1(-x), \eta_1(-x))$ is also a positive eigenvector. From the simplicity of eigenvector in Lemma 2.4, it is necessary that $(\xi_1(-x), \eta_1(-x)) = (\xi_1(x), \eta_1(x))$ for $x \in I$. This implies that (ξ_1, η_1) is symmetric with respect to 0, thus (μ_1, ξ_1, η_1) satisfies (2.26). Therefore the principal eigenvalues of (2.21) and (2.26) are equivalent. In particular, a solution (u_1, u_2) of (2.2) is stable if the principal eigenvalue μ_1 of (2.26) is positive.

The stability of a positive crossing solution is known if (f, g) is sublinear or superlinear. The following lemma is from Theorem 2.3 and Theorem 2.4 of [25].

Lemma 2.5. *Let (u_1, u_2) be a solution of (2.2) such that $u_1'(R) < 0$ and $u_2'(R) < 0$. Assume that (f, g) is cooperative as defined in (f2).*

- (1) *If (f, g) is sublinear, then the principal eigenvalue μ_1 of (2.26) is positive, i.e., (u_1, u_2) is stable;*
- (2) *If (f, g) is superlinear, then the principal eigenvalue μ_1 of (2.26) is negative, i.e., (u_1, u_2) is unstable.*

Next we observe a connection between the principal eigenvalue of (2.26) (or equivalently (2.21)) and two scalar eigenvalue problems. Consider the eigenvalue problems

$$(2.27) \quad \begin{cases} \xi_3'' + f_1 \xi_3 = -\kappa \xi_3, & 0 < x < R, \\ \xi_3'(0) = 0, \quad \xi_3(R) = 0, \end{cases}$$

and

$$(2.28) \quad \begin{cases} \eta_3'' + g_2 \eta_3 = -\nu \eta_3, & 0 < x < R, \\ \eta_3'(0) = 0, \quad \eta_3(R) = 0. \end{cases}$$

Then we have the relation:

Proposition 2.6. *Let (u_1, u_2) be a solution of (2.2) such that $u_1'(R) < 0$ and $u_2'(R) < 0$, and assume that (f, g) is cooperative as defined in (f2). Let μ_1 be the principal eigenvalue of (2.26), let κ_1 and ν_1 be the principal eigenvalues of (2.27) and (2.28) respectively, and let (ϕ_3, ψ_3) be the solution of (2.8). Then*

$$(2.29) \quad \mu_1 < \min\{\kappa_1, \nu_1\}.$$

In particular, if one of ϕ_3 and ψ_3 changes sign in $(0, R)$, then (u_1, u_2) is unstable.

Proof. Let $L_1(\phi) = -\phi'' - f_1\phi - \mu_1\phi$ for $\phi \in H_0^1(I)$. Then $L_1(\xi_1) = f_2\eta_1 \geq (\neq)0$ for $x \in I$, and $\xi_1 = 0$ for $x = \pm R$. Hence from the maximum principle (part 5 of Lemma 2.4 for L_1), the principal eigenvalue $\mu_1(L_1) = \kappa_1 - \mu_1 > 0$. Same proof can be applied to $L_2(\phi) = -\phi'' - g_2\phi - \mu_1\phi$ for $\phi \in H_0^1(I)$ to prove $\mu_1 < \nu_1$. It is well-known that κ_1 (or ν_1) is positive if and only if ϕ_3 (or ψ_3) is positive in $[0, R]$ from Sturm comparison lemma. Thus if one of ϕ_3 and ψ_3 changes sign in $(0, R)$, then $\mu_1 < \min\{\kappa_1, \nu_1\} < 0$. \square

Remark 2.7.

- (1) Some stability results similar to Lemma 2.5 for a positive solution $u(x)$ of the scalar elliptic equation

$$(2.30) \quad \Delta u + h(u) = 0, \quad x \in \Omega, \quad u(x) = 0, \quad x \in \partial\Omega,$$

were proved in [54] for the general bounded spatial domain $\Omega \subseteq \mathbb{R}^n$. If $h(u)$ is sublinear, that is $h'(u)u - h(u) \leq 0$ for $u \geq 0$, then u is stable; and if $h(u)$ is superlinear, that is $h'(u)u - h(u) \geq 0$ for $u \geq 0$, then u is unstable.

- (2) If (f, g) is weakly sublinear, and (u_1, u_2) is a positive solution of (1.3), then both ϕ_3 and ψ_3 are positive on $[0, R]$. But (u_1, u_2) may not be stable with respect to (1.3). For example, consider the boundary value problem

$$(2.31) \quad \begin{cases} u_1'' + u_1 - u_1^3 + u_1 u_2^2 = 0, & x \in (-\pi/2, \pi/2), \\ u_2'' + u_2 - u_2^3 + u_1^2 u_2 = 0, & x \in (-\pi/2, \pi/2), \\ u_1(x) > 0, \quad u_2(x) > 0, & x \in (-\pi/2, \pi/2), \\ u_1(\pm\pi/2) = u_2(\pm\pi/2) = 0. \end{cases}$$

It is easy to verify that $(u_1(x), u_2(x)) = (\cos(x), \cos(x))$ is a positive solution to (2.31). This solution is neutrally stable as $\mu_1 = 0$ and $(\xi_1, \eta_1) = (\cos(x), \cos(x))$. Since $f_1 u_1 - f = -2u_1^3 < 0$ and $g_2 u_2 - g = -2u_2^3 < 0$, then (f, g) is weakly sublinear. But (f, g) is not sublinear since $f_1 u_1 + f_2 u_2 - f = 2u_1(u_2^2 - u_1^2)$ and $g_1 u_1 + g_2 u_2 - g = 2u_2(u_1^2 - u_2^2)$.

2.3. Nondegeneracy. With the oscillatory properties of (ϕ_i, ψ_i) ($i = 1, 2$) proved in Lemma 2.2 and Corollary 2.3, we consider the solution (A_c, B_c) of the following initial value problem:

$$(2.32) \quad \begin{cases} A_c'' + f_1 A_c + f_2 B_c = 0, & 0 < x < R, \\ B_c'' + g_1 A_c + g_2 B_c = 0, & 0 < x < R, \\ A_c(0) = 1, \quad A_c'(0) = 0, \\ B_c(0) = c > 0, \quad B_c'(0) = 0, \end{cases}$$

for $c \geq 0$. Clearly $(A_c, B_c) = (\phi_1, \psi_1) + c(\phi_2, \psi_2)$. If $(u_1(x; \alpha, \beta), u_2(x; \alpha, \beta))$ is a crossing solution of (2.2), then the behavior of the solution (A_c, B_c) roughly shows the nodal behavior of $(u_1(x; \alpha + \varepsilon, \beta + c\varepsilon) - u_1(x; \alpha, \beta), u_2(x; \alpha + \varepsilon, \beta + c\varepsilon) - u_2(x; \alpha, \beta))$, thus (A_c, B_c) can be viewed as the directional derivative of (u_1, u_2) along the direction $(1, c)$.

We first show a general result on the connection between the stability of a solution and the variational equation (2.32).

Proposition 2.8. Let (u_1, u_2) be a solution of (2.2) such that $u_1'(R) < 0$ and $u_2'(R) < 0$ and let (A_c, B_c) be defined as in (2.32). Assume that (f, g) is cooperative as defined in (f2).

- (1) (u_1, u_2) is stable if and only if for some $c > 0$, $A_c(x) > 0$, $B_c(x) > 0$ in $(0, R]$;
- (2) (u_1, u_2) is neutrally stable if and only if for some $c > 0$, $A_c(x) > 0$ and $B_c(x) > 0$ in $(0, R)$ and $A_c(R) = B_c(R) = 0$;
- (3) (u_1, u_2) is unstable if and only if for all $c \geq 0$, at least one of $A_c(x)$ or $B_c(x)$ is not positive in $(0, R)$.

Proof. For part 1, if for some $c > 0$, $A_c(x) > 0$ and $B_c(x) > 0$ in $(0, R]$, then we extend (A_c, B_c) to $[-R, R]$. Then $\mu_1 > 0$ follows from part 5 of Lemma 2.4. On

the other hand, if (u_1, u_2) is stable, then (2.21) has a positive principal eigenvalue $\mu_1(R) > 0$ with a positive eigenvector (ξ_1, η_1) . From the uniqueness of principal eigenvector (Lemma 2.4 part 2) and the fact that (u_1, u_2) is symmetric with respect to $x = 0$, then (ξ_1, η_1) is also symmetric with respect to $x = 0$, hence (μ_1, ξ_1, η_1) necessarily satisfies (2.26). Recall that we have extended the definition of (f, g) to \mathbb{R}^2 so it is C^1 , thus we can also extend $f_i(u_1(x), u_2(x)), g_i(u_1(x), u_2(x)), i = 1, 2$, to $x \in \mathbb{R}^2$. Consider the eigenvalue problem:

$$(2.33) \quad \begin{cases} \xi'' + f_1\xi + f_2\eta = -\mu\xi, & x \in I_r, \\ \eta'' + g_1\xi + g_2\eta = -\mu\eta, & x \in I_r, \\ \xi(x) = 0, \quad \eta(x) = 0, & x \in \partial I_r, \end{cases}$$

where $I_r = (-r, r)$ for $r > 0$. It is well known that the principal eigenvalue $\mu_1(r)$ is continuous with respect to r . Moreover, if (f, g) is cooperative, then $\mu_1(r)$ is non-increasing with respect to r by (2.25). We claim that there exists some $c > 0$ such that $A_c(x) > 0, B_c(x) > 0$ in $(0, R]$. On the contrary, suppose for any $c > 0, A_c(x)$ or $B_c(x)$ changes sign at least once in $(0, R]$. Then the graphs of $c_1(x)$ and $c_2(x)$ must intersect in $(0, R]$. Let x_* be the smallest $x > 0$ such that $c_1(x) = c_2(x) = c > 0$. We obtain that the principal eigenvalue of (2.33) is zero, *i.e.* $\mu_1(x_*) = 0$. It implies that $\mu_1(R) \leq 0$ since $x_* \leq R$. This contradicts with $\mu_1(R) > 0$.

Part 2 follows easily from part 2 of Lemma 2.4 and the definition of neutral stability. Part 3 also follows easily from part 1 and 2 as (u_1, u_2) is unstable if it is not stable nor neutrally stable. \square

Proposition 2.8 shows that the oscillatory property of functions A_c and B_c is closely related to the stability of a positive solution (u_1, u_2) of (2.2). The sign-changing of the functions A_c and B_c can be tracked by the functions

$$(2.34) \quad c_1(x) = -\frac{\phi_1(x)}{\phi_2(x)}, \quad 0 < x \leq R, \quad c_2(x) = -\frac{\psi_1(x)}{\psi_2(x)}, \quad 0 < x \leq R, \quad \psi_2(x) \neq 0.$$

From Lemma 2.1, $c_1(x)$ is well-defined for all $x \in (0, R]$; $c_2(x)$ is well-defined for all $x \in (0, R]$ if $\psi_2(x) > 0$ for $x \in (0, R]$, and $c_2(x)$ has a vertical asymptote at some $x_* \in (0, R)$ if $\psi_2(x_*) = 0$ for some $x_* \in (0, R)$. For a given $c > 0$, the roots of $c_1(x) = c$ and $c_2(x) = c$ give the zeros of the functions $A_c(x)$ and $B_c(x)$.

First we show that the stability/instability result for the sublinear/superlinear cases shown in Lemma 2.5 can be rephrased using $c_1(x)$ and $c_2(x)$ as follows.

Lemma 2.9. *Let (u_1, u_2) be a solution of (2.2) such that $u_1'(R) < 0$ and $u_2'(R) < 0$, and let $c_i(x)$ ($i = 1, 2$) be defined as in (2.34). Assume that (f, g) is cooperative as defined in (f2).*

- (1) *If (f, g) is sublinear, then $c_1(x)$ is strictly decreasing and $c_2(x)$ is strictly increasing for $x \in (0, R)$, and $c_1(R) > c_2(R)$, *i.e.* the graphs of $c_1(x)$ and $c_2(x)$ do not intersect in $(0, R]$;*
- (2) *If (f, g) is superlinear, then there exists $x^* \in (0, R)$ such that the graphs of $c_1(x)$ and $c_2(x)$ intersect at $x = x^*$, and $c_1(x)$ is strictly decreasing and $c_2(x)$ is strictly increasing for $x \in (0, x^*)$.*

Proof. 1. If (f, g) is sublinear, then $\phi_1(x) > 0$ and $\psi_2(x) > 0$ for all $x \in [0, R]$ by Corollary 2.3. Thus $c_1(x), c_2(x) > 0$ for $x \in (0, R]$. We prove that $c_1(x) > c_2(x)$ for

all $x \in [0, R]$. When $x > 0$ is small, we have $c_1(x) > c_2(x)$ since $\lim_{x \rightarrow 0^+} c_1(x) = \infty$ and $c_2(0) = 0$. Let x_* be the smallest $x > 0$ such that $c_1(x) = c_2(x) = c^* > 0$. From the definition of $c_i(x)$, we have

$$(2.35) \quad \begin{aligned} c_1'(x) &= \frac{\phi_1 \phi_2' - \phi_2 \phi_1'}{\phi_2^2} = \frac{\int_0^x f_2(\psi_1 \phi_2 - \psi_2 \phi_1) ds}{\phi_2^2} = \frac{\int_0^x f_2 \phi_2 \psi_2 (c_1 - c_2) ds}{\phi_2^2}, \\ c_2'(x) &= \frac{\psi_1 \psi_2' - \psi_2 \psi_1'}{\psi_2^2} = \frac{\int_0^x g_1(\psi_2 \phi_1 - \psi_1 \phi_2) ds}{\psi_2^2} = \frac{\int_0^x g_1 \phi_2 \psi_2 (c_2 - c_1) ds}{\psi_2^2}. \end{aligned}$$

Since $c_1(x) - c_2(x) > 0$ for $x \in (0, x_*)$, then $c_1'(x) < 0$, $c_2'(x) > 0$, and $c_1(x) > c^* > c_2(x)$ for $x \in (0, x_*)$. This also implies that for $c = c^*$, $A_{c^*}(x) > 0$ and $B_{c^*}(x) > 0$ for $x \in [0, x_*)$ and $A_{c^*}(x_*) = B_{c^*}(x_*) = 0$.

Consider an eigenvalue problem

$$(2.36) \quad \begin{cases} \xi'' + f_1 \xi + f_2 \eta = -\mu \xi, & x \in (0, x_*), \\ \eta'' + g_1 \xi + g_2 \eta = -\mu \eta, & x \in (0, x_*), \\ \xi'(0) = \xi(x_*) = 0, \quad \eta'(0) = \eta(x_*) = 0. \end{cases}$$

From the above discussion and Lemma 2.4, we know that the principal eigenvalue of (2.36) is zero, *i.e.* $\mu_1(x_*) = 0$ and $(A_{c^*}(x), B_{c^*}(x))$ with $x \in (0, x_*)$ is the corresponding positive eigenfunction. On the other hand, by (2.25), we have $\mu_1(R) \leq \mu_1(x_*) = 0$ since $x_* \leq R$. But since (f, g) is sublinear, then $\mu_1(R) > 0$ from part 1 of Lemma 2.5. That is a contradiction. Hence if (f, g) is sublinear, then $c_1(x) > c_2(x)$ for all $x \in [0, R]$. It implies that for any $c \in (c_2(R), c_1(R))$, $A_c(x) > 0$ and $B_c(x) > 0$ in $(0, R]$.

2. Next we assume that (f, g) is superlinear. We prove that $c_1(x)$ and $c_2(x)$ must intersect at some $x^* \in (0, R)$. Suppose not, then $c_1(x) > c_2(x)$ for $x \in (0, R)$ and $c_1(R) \geq c_2(R)$. Let $c = c_1(R)$, then $A_c(x) > 0$ and $B_c(x) > 0$ in $(0, R)$. If $c_1(R) > c_2(R)$, then $B_c(R) > 0$ so $\mu_1(R) > 0$ from part 5 of Lemma 2.4. If $c_1(R) = c_2(R)$, then $\mu_1(R) = 0$. Therefore we get $\mu_1(R) \geq 0$, which contradicts with part 2 of Lemma 2.5. \square

To prove the non-degeneracy of positive solution for superlinear problem, we would need the information of $c_1(x)$ and $c_2(x)$ beyond the intersection point shown in Lemma 2.9. While in general it is difficult to classify such behavior, in the following key lemma, we prove that the graphs of $c_1(x)$ and $c_2(x)$ cannot intersect again if addition structure is imposed on the vector field (f, g) : (i) it is weakly sublinear; and (ii) it is variational (gradient or Hamiltonian).

Lemma 2.10. *Let (u_1, u_2) be a solution of (2.2) such that $u_1'(R) < 0$ and $u_2'(R) < 0$, and let $c_i(x)$ ($i = 1, 2$) be defined as in (2.34). Assume that (f, g) is cooperative as defined in (f2). If (f, g) is superlinear and weakly sublinear, and (2.2) is a Hamiltonian system or a gradient system, then the graphs of $c_1(x)$ and $c_2(x)$ intersect only at $x = x^*$ in $(0, R]$, each of $c_1(x)$ and $c_2(x)$ has at most one critical point in $(0, R]$, and $c_2(x) > c_1(x)$ for $x \in (x^*, R]$.*

Proof. We assume that (f, g) is superlinear and weakly sublinear. From part 2 of Lemma 2.9, the graphs of $c_1(x)$ and $c_2(x)$ intersect at some $x^* \in (0, R)$. We prove that the graphs of $c_1(x)$ and $c_2(x)$ do not intersect in $(x^*, R]$. Since (f, g) is weakly sublinear, then $\phi_1(x) > 0$ and $\psi_2(x) > 0$ for all $x \in [0, R]$ by Corollary 2.3. From (2.35), $c_1'(x) < 0$ and $c_2'(x) > 0$ for $x \in (0, x^*]$, and thus there exists small $\epsilon > 0$

such that $c_1(x) < c_2(x)$ for $x \in (x^*, x^* + \epsilon]$. Suppose that the graphs of $c_1(x)$ and $c_2(x)$ have another intersection point. Let x^{**} be the smallest $x \in (x^*, R]$ such that $c_1(x) = c_2(x) = c^{**} > 0$. Then $c_1(x) < c_2(x)$ for $x \in (x^*, x^{**})$.

Note that

$$(2.37) \quad \begin{aligned} c_1''(x) &= \frac{f_2 \phi_2 \psi_2 (c_1 - c_2) \phi_2^2 - 2 \phi_2 \phi_2' \int_0^x f_2 \phi_2 \psi_2 (c_1 - c_2) ds}{\phi_2^4}, \\ c_2''(x) &= \frac{g_1 \phi_2 \psi_2 (c_2 - c_1) \psi_2^2 - 2 \psi_2 \psi_2' \int_0^x g_1 \phi_2 \psi_2 (c_2 - c_1) ds}{\psi_2^4}. \end{aligned}$$

If there exists $x_1 \in (x^*, x^{**})$ such that $c_1'(x_1) = 0$, then

$$c_1''(x_1) = \left. \frac{f_2 \phi_2 \psi_2 (c_1 - c_2)}{\phi_2^2} \right|_{x=x_1} > 0,$$

since $c_1(x_1) < c_2(x_1)$ and $f_2(x) > 0$, $\phi_2(x) < 0$, $\psi_2(x) > 0$ for $x \in (0, R]$. Similarly, if there exists $x_2 \in (x^*, x^{**})$ such that $c_2'(x_2) = 0$, then

$$c_2''(x_2) = \left. \frac{g_1 \phi_2 \psi_2 (c_2 - c_1)}{\psi_2^2} \right|_{x=x_2} < 0.$$

This combining with $c_1'(x) < 0$ and $c_2'(x) > 0$ in $(0, x^*]$ implies that $c_1(x)$ has at most one critical point which is a local minimum, and $c_2(x)$ has at most one critical point which is a local maximum. In particular, the horizontal line $c = c^{**}$ intersects each of $c = c_1(x)$ and $c = c_2(x)$ at most once for $x \in [0, x^{**})$, or equivalently, each of $A_{c^{**}}(x)$ and $B_{c^{**}}(x)$ changes sign in $(0, x^{**})$ at most once. Then there are following three possible cases:

- (i) Both of $A_{c^{**}}(x)$ and $B_{c^{**}}(x)$ change sign in $(0, x^{**})$ exactly once.
- (ii) $A_{c^{**}}(x)$ changes sign in $(0, x^{**})$ exactly once, and $B_{c^{**}}(x)$ does not change sign in $(0, x^{**})$.
- (iii) $B_{c^{**}}(x)$ changes sign in $(0, x^{**})$ exactly once, and $A_{c^{**}}(x)$ does not change sign in $(0, x^{**})$.

In the following, we discuss these three cases when one of two types of additional variational structure (Hamiltonian or gradient) is imposed on system (2.2).

Type I: First we suppose (2.2) is a Hamiltonian system. Then

$$(2.38) \quad f_1(u_1, u_2) \equiv g_2(u_1, u_2), \quad \text{for } (u_1, u_2) \in \mathbb{R}_+^2.$$

If case (i) occurs, then

$$(2.39) \quad A_{c^{**}}(x^{**}) = B_{c^{**}}(x^{**}) = 0, \quad A'_{c^{**}}(x^{**}) \geq 0, \quad B'_{c^{**}}(x^{**}) \geq 0.$$

Using the equation of A_c with $c = c^{**}$ and the equation of u_2' , we obtain

$$(2.40) \quad A''_{c^{**}} u_2' - (u_2')'' A_{c^{**}} + f_2 B_{c^{**}} u_2' - g_1 u_1' A_{c^{**}} = 0,$$

since (2.38) holds. Similarly we have

$$(2.41) \quad B''_{c^{**}} u_1' - (u_1')'' B_{c^{**}} + g_1 u_1' A_{c^{**}} - f_2 u_2' B_{c^{**}} = 0.$$

Adding (2.40) and (2.41), we get

$$(2.42) \quad A''_{c^{**}} u_2' - (u_2')'' A_{c^{**}} + B''_{c^{**}} u_1' - (u_1')'' B_{c^{**}} = 0.$$

Define a function

$$P(x) = A'_{c^{**}}(x) u_2'(x) - u_2''(x) A_{c^{**}}(x) + B'_{c^{**}}(x) u_1'(x) - u_1''(x) B_{c^{**}}(x), \quad x \in [0, R].$$

Then (2.42) implies that $P'(x) \equiv 0$ for $x \in (0, R)$. Hence, for $x \in [0, R]$,

$$(2.43) \quad P(x) \equiv P(0) = -u_2''(0)A_{c^{**}}(0) - u_1''(0)B_{c^{**}}(0) = g(\alpha, \beta) + f(\alpha, \beta)c^{**} > 0.$$

However, from (2.39), we have

$$(2.44) \quad P(x^{**}) = A'_{c^{**}}(x^{**})u_2'(x^{**}) + B'_{c^{**}}(x^{**})u_1'(x^{**}) \leq 0,$$

which is a contradiction with (2.43).

If case (ii) occurs, suppose the unique zero of $A_{c^{**}}$ in $(0, x^{**})$ is x_1 , then

$$(2.45) \quad A_{c^{**}}(x_1) = A_{c^{**}}(x^{**}) = B_{c^{**}}(x^{**}) = 0, \quad A_{c^{**}}(x) < 0, \quad B_{c^{**}}(x) > 0 \text{ in } (x_1, x^{**}),$$

and

$$(2.46) \quad A'_{c^{**}}(x_1) \leq 0, \quad A'_{c^{**}}(x^{**}) \geq 0.$$

Then multiplying the equation of $A_{c^{**}}$ by u_1' , multiplying the equation of u_1' in (2.9) by $A_{c^{**}}$, subtracting and integrating on (x_1, x^{**}) , we obtain

$$(2.47) \quad A'_{c^{**}}(x^{**})u_1'(x^{**}) - A'_{c^{**}}(x_1)u_1'(x_1) = \int_{x_1}^{x^{**}} f_2 u_2'(A_{c^{**}} - B_{c^{**}}) dx.$$

The left hand side of (2.47) is non-positive since $u_1'(x) < 0$ in $(0, R)$ and (2.46) holds, while the right hand side of (2.47) is positive since $f_2 > 0$, $u_2' < 0$ in $(0, R)$ and (2.45) holds. That is a contradiction. If case (iii) occurs, we can derive a similar contradiction as case (ii).

Summarizing discussion above, we have proved that if (f, g) is a Hamiltonian system, then $c_1(x) < c_2(x)$ for all $x \in (x^*, R]$, $c_1(x)$ has at most one critical point which is a local minimum, and $c_2(x)$ has at most one critical point which is a local maximum for $x \in (0, R)$.

Type II: Secondly, we suppose (2.2) is a gradient system. Then

$$(2.48) \quad f_2(u_1, u_2) \equiv g_1(u_1, u_2), \quad \text{for } (u_1, u_2) \in \mathbb{R}_+^2.$$

If case (i) occurs, then again (2.39) holds. Using the equation of A_c with $c = c^{**}$ and the equation of u_1' , we obtain

$$(2.49) \quad A_{c^{**}}'' u_1' - (u_1')'' A_{c^{**}} + f_2 B_{c^{**}} u_1' - f_2 u_2' A_{c^{**}} = 0,$$

since (2.48) holds. Similarly we have

$$(2.50) \quad B_{c^{**}}'' u_2' - (u_2')'' B_{c^{**}} + g_1 u_2' A_{c^{**}} - g_1 u_2' B_{c^{**}} = 0.$$

Adding (2.49) and (2.50), we get that

$$(2.51) \quad A_{c^{**}}'' u_1' - (u_1')'' A_{c^{**}} + B_{c^{**}}'' u_2' - (u_2')'' B_{c^{**}} = 0.$$

Define a function

$$Q(x) = A'_{c^{**}}(x)u_1'(x) - u_1''(x)A_{c^{**}}(x) + B'_{c^{**}}(x)u_2'(x) - u_2''(x)B_{c^{**}}(x), \quad x \in [0, R].$$

Then we have $Q'(x) \equiv 0$ for $x \in (0, R)$ from (2.51), which implies that for any $x \in [0, R]$,

$$(2.52) \quad Q(x) \equiv Q(0) = -u_1''(0)A_{c^{**}}(0) - u_2''(0)B_{c^{**}}(0) = f(\alpha, \beta) + g(\alpha, \beta)c^{**} > 0.$$

On the other hand, from (2.39), we have

$$(2.53) \quad Q(x_*) = A'_{c^{**}}(x_*)u_1'(x_*) + B'_{c^{**}}(x_*)u_2'(x_*) \leq 0,$$

that is a contradiction. The proofs for case (ii) and (iii) are the same as the ones in the proof of Hamiltonian case. This completes the proof for gradient system case. \square

In the above proof, we show that the superlinearity of the vector field (f, g) ensures that $c_1(x)$ and $c_2(x)$ intersect at least once for $x \in (0, R]$, while the weakly sublinearity and variational structure (gradient or Hamiltonian) of (f, g) ensure that $c_1(x)$ and $c_2(x)$ intersect only once for $x \in (0, R]$. Both of these two aspects guarantee the non-degeneracy of a positive solution of problem (2.2). In Section 3 we will give two examples of vector fields with variational structure (gradient or Hamiltonian) and satisfying simultaneously the superlinear and weakly sublinear conditions.

Define

$$(2.54) \quad c_1 = c_1(R) = \begin{cases} -\frac{\phi_1(R)}{\phi_2(R)}, & \text{if } \phi_1(R) > 0, \\ 0, & \text{if } \phi_1(R) \leq 0; \end{cases} \quad c_2 = c_2(R) = \begin{cases} -\frac{\psi_1(R)}{\psi_2(R)}, & \text{if } \psi_2(R) > 0, \\ \infty, & \text{if } \psi_2(R) \leq 0. \end{cases}$$

The properties of $c_1(x)$ and $c_2(x)$ in Lemma 2.9 imply the following corollary, which is the key for obtaining the uniqueness of positive solution in the next section.

Corollary 2.11. Let (u_1, u_2) be a solution of (2.2) such that $u_1'(R) < 0$ and $u_2'(R) < 0$. Assume that (2.3) is a Hamiltonian system or is a gradient system, and (f, g) is cooperative as defined in (f2). Then

- (1) If (f, g) is sublinear, then $c_1 > c_2$, and for any $c_2 < c < c_1$, each of $A_c(x)$ and $B_c(x)$ is positive in $(0, R]$;
- (2) If (f, g) is superlinear and weakly sublinear, then $c_1 < c_2$, and for any $c_1 < c < c_2$, each of $A_c(x)$ and $B_c(x)$ changes sign exactly once in $(0, R)$ and $A_c(R) < 0, B_c(R) < 0$. Moreover for any $c \geq c_2$ or $c \leq c_1$, $A_c(R)B_c(R) \leq 0$.

Proof. 1. If (f, g) is sublinear, then $c_1 = c_1(R) > c_2(R) = c_2$ from part 1 of Lemma 2.9. Clearly for $c \in (c_2, c_1)$, each of A_c and B_c is positive from definition.

2. If (f, g) is superlinear and weakly sublinear, then $c_1 = c_1(R) < c_2(R) = c_2$ from parts 2 and 3 of Lemma 2.9. Let $c_m = \min_{x \in (0, R]} c_1(x)$ and $c_M = \max_{x \in (0, R]} c_2(x)$. Then

from parts 2 and 3 of Lemma 2.9, we have $c_m \leq c_1 < c_2 \leq c_M$. If $c \in (c_2, c_1)$, then each of $c_1(x)$ and $c_2(x)$ equals to c exactly once for $x \in (0, R]$, hence each of $A_c(x)$ and $B_c(x)$ changes sign exactly once in $(0, R)$. If $c \geq c_2$, then we always have A_c changes sign exactly once and $A_c(R) < 0$. For B_c , there are several cases: (i) if $c > c_M$, then B_c is positive; (ii) if $c = c_M$, then B_c is positive except at one point; (iii) if $c_2 < c < c_M$, then B_c changes sign exactly twice. In all three cases, we have $B_c(R) > 0$. Thus $A_c(R)B_c(R) < 0$ if $c > c_2$. When $c = c_2$, we have $B_c(R) = 0$ hence $A_c(R)B_c(R) = 0$. Similarly we can also show that for any $c \leq c_1$, $A_c(R)B_c(R) \leq 0$. \square

For the sublinear case, the functions $c_1(x)$ and $c_2(x)$ defined in (2.34) have been used in [13, 14] for special Hamiltonian nonlinearity but for the radially symmetric solutions in higher dimension, and again one can show that the graphs of $c_1(x)$ and $c_2(x)$ do not intersect. Here we prove such result hold for any sublinear system in Lemma 2.9. The results in Lemma 2.10 and Corollary 2.11 appear to be the

first ones considering $c_1(x)$ and $c_2(x)$ intersecting, and we obtain the nondegeneracy results under three conditions on the vector field: (i) superlinear; (ii) weakly sublinear; and (iii) Hamiltonian or gradient. Applications of these results will be given in Section 3.

3. UNIQUENESS FOR WEAKLY SUBLINEAR AND SUPERLINEAR SYSTEMS

3.1. A Hamiltonian Schrödinger system. In this subsection we consider (1.2) and (1.3) with the following Hamiltonian functional:

$$(3.1) \quad H(u_1, u_2) = -u_1 u_2 + H_1(u_1) + H_2(u_2),$$

where for $i = 1, 2$, $H_i : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is C^2 function satisfying $H_i(0) = 0$. Let $h_i(u_i) = H_i'(u_i)$. Then the corresponding elliptic system on a bounded interval is

$$(3.2) \quad \begin{cases} u_1'' - u_1 + h_2(u_2) = 0, & x \in (0, R), \\ u_2'' - u_2 + h_1(u_1) = 0, & x \in (0, R), \\ u_1(x) > 0, \quad u_2(x) > 0, & x \in (0, R), \\ u_1'(0) = u_2'(0) = 0, \quad u_1(R) = u_2(R) = 0, \end{cases}$$

and the ground state solutions satisfy

$$(3.3) \quad \begin{cases} u_1'' - u_1 + h_2(u_2) = 0, & x \in (0, \infty), \\ u_2'' - u_2 + h_1(u_1) = 0, & x \in (0, \infty), \\ u_1(x) > 0, \quad u_2(x) > 0, \quad u_1'(x) < 0, \quad u_2'(x) < 0, & x \in (0, \infty), \\ u_1'(0) = u_2'(0) = 0. \end{cases}$$

In this subsection we assume that for $i = 1, 2$,

$$(3.4) \quad h_i(0) = 0, \quad h_i'(u_i) > 0, \quad \text{and} \quad h_i'(u_i)u_i - h_i(u_i) > 0 \quad \text{for} \quad u_i > 0;$$

and

$$(3.5) \quad h_i'(0) = 0, \quad \lim_{u_i \rightarrow \infty} h_i'(u_i) = \infty.$$

Notice that (3.4) implies that (f, g) is superlinear but not strongly superlinear. Also (f, g) is weakly sublinear.

We consider the initial value problem

$$(3.6) \quad \begin{cases} u_1'' - u_1 + h_2(u_2) = 0, & x > 0, \\ u_2'' - u_2 + h_1(u_1) = 0, & x > 0, \\ u_1'(0) = u_2'(0) = 0, \\ u_1(0) = \alpha > 0, \quad u_2(0) = \beta > 0. \end{cases}$$

Define

$$(3.7) \quad f(u_1, u_2) = -u_1 + h_2(u_2), \quad \text{and} \quad g(u_1, u_2) = -u_2 + h_1(u_1).$$

According to the signs of f and g , we define the following regions in \mathbb{R}_+^2 :

$$(3.8) \quad \begin{aligned} I &= \{(u_1, u_2) \in \mathbb{R}_+^2 : f(u_1, u_2) > 0, g(u_1, u_2) > 0\}, \\ II &= \{(u_1, u_2) \in \mathbb{R}_+^2 : f(u_1, u_2) < 0, g(u_1, u_2) < 0\}, \\ III &= \{(u_1, u_2) \in \mathbb{R}_+^2 : f(u_1, u_2) < 0, g(u_1, u_2) > 0\}, \\ IV &= \{(u_1, u_2) \in \mathbb{R}_+^2 : f(u_1, u_2) > 0, g(u_1, u_2) < 0\}. \end{aligned}$$

Since we assume that h_i satisfies (3.4) and (3.5), then the curves $f(u_1, u_2) = 0$ and $g(u_1, u_2) = 0$ are monotone ones, and they have a unique intersection point (u_1^*, u_2^*) (see Figure 1).

For $(\alpha, \beta) \in II \cup III \cup IV$, $u_1' > 0$ or $u_2' > 0$ in $(0, \delta)$, hence it cannot be a solution of (3.2). For $(\alpha, \beta) \in I$, $u_1' < 0$ and $u_2' < 0$ in $(0, \delta)$. We recall $R = R(\alpha, \beta)$ to be the right endpoint of the maximal interval $(0, R(\alpha, \beta))$ so that $u_i(x) > 0$ and $u_i'(x) < 0$ in $(0, R(\alpha, \beta))$, $i = 1, 2$. We partition I into the following classes:

$$\begin{aligned}
(3.9) \quad \mathcal{B} &= \{(\alpha, \beta) \in I : R < \infty, u_1(R) = 0, u_1'(R) < 0, u_2(R) > 0, u_2'(R) < 0\}, \\
\mathcal{G} &= \{(\alpha, \beta) \in I : R < \infty, u_1(R) > 0, u_1'(R) = 0, u_2(R) > 0, u_2'(R) < 0\}, \\
\mathcal{R} &= \{(\alpha, \beta) \in I : R < \infty, u_1(R) > 0, u_1'(R) < 0, u_2(R) = 0, u_2'(R) < 0\}, \\
\mathcal{Y} &= \{(\alpha, \beta) \in I : R < \infty, u_1(R) > 0, u_1'(R) < 0, u_2(R) > 0, u_2'(R) = 0\}, \\
\mathcal{S} &= \{(\alpha, \beta) \in I : R < \infty, u_1(R) = 0, u_1'(R) < 0, u_2(R) = 0, u_2'(R) < 0\}, \\
\mathcal{Q} &= \{(\alpha, \beta) \in I : R = \infty, \lim_{x \rightarrow \infty} u_1(x) = \lim_{x \rightarrow \infty} u_2(x) = 0\}, \\
\mathcal{P} &= I \setminus (\mathcal{B} \cup \mathcal{G} \cup \mathcal{R} \cup \mathcal{Y} \cup \mathcal{S} \cup \mathcal{Q}).
\end{aligned}$$

It is clear that if $(\alpha, \beta) \in \mathcal{S}$, then the corresponding solution (u_1, u_2) is a solution of (3.2), while each element in \mathcal{Q} defines a ground state solution in the whole space. The elements in \mathcal{S} and \mathcal{Q} can be characterized as follows.

Lemma 3.1. *Consider equation (3.6), and let $H(u_1, u_2)$ be defined as in (3.1).*

- (1) *If $(\alpha, \beta) \in \mathcal{S}$, then $H(\alpha, \beta) > 0$, $u_1'(R) < 0$ and $u_2'(R) < 0$.*
- (2) *If $(\alpha, \beta) \in \mathcal{Q}$, then $H(\alpha, \beta) = 0$, and $u_1(x), u_2(x), u_1'(x), u_2'(x) \rightarrow 0$ as $x \rightarrow \infty$.*

Proof. For part 1, if $(\alpha, \beta) \in \mathcal{S}$, then $(u_1(x), u_2(x))$ satisfies

$$\begin{aligned}
(3.10) \quad u_1'' - u_1 &= -h_2(u_2) < 0, \quad u_2'' - u_2 = -h_1(u_1) < 0, \quad x \in (-R, R), \\
u_1 = u_2 &= 0, \quad |x| = R.
\end{aligned}$$

The maximum principle and the Hopf boundary lemma hold, hence $u_1'(R) < 0$ and $u_2'(R) < 0$. Recall $H_0(x)$ defined in (1.14). Since $H_0(x) \equiv H_0(0) = H(\alpha, \beta)$, then $H(\alpha, \beta) = H_0(R) = u_1'(R)u_2'(R) > 0$.

For part 2, assume that $(\alpha, \beta) \in \mathcal{Q}$. It is clear that $u_i'(x), u_i''(x) \rightarrow 0$, and $(u_1(x), u_2(x)) \rightarrow (a, b) \in \mathbb{R}_+^2$ as $x \rightarrow \infty$. Since $u_i''(x) \rightarrow 0$ as $x \rightarrow \infty$, then $f(a, b) = g(a, b) = 0$. Thus (a, b) is either $(0, 0)$ or (u_1^*, u_2^*) . We claim that the latter case is not possible. In fact, from $H_0(x) \equiv H_0(0) = H(\alpha, \beta)$, and $u_i'(x) \rightarrow 0$ as $x \rightarrow \infty$, we conclude that $H(a, b) = H(\alpha, \beta)$. But (u_1^*, u_2^*) is the global minimum of the energy functional H , and for any $(\alpha, \beta) \in \mathbb{R}_+^2$, $(\alpha, \beta) \neq (u_1^*, u_2^*)$, $H(\alpha, \beta) > H(u_1^*, u_2^*)$. Thus $(\alpha, \beta) = (u_1^*, u_2^*)$ if $(a, b) = (u_1^*, u_2^*)$. That is a contradiction. Hence $(a, b) = (0, 0)$. And $0 = \lim_{x \rightarrow \infty} H_0(x) \equiv H_0(0)$ implies that $H(\alpha, \beta) = H_0(0) = 0$. \square

For the purpose of identifying the sets \mathcal{S} and \mathcal{Q} , we will use a more coarse partition of \mathbb{R}_+^2 than the one given in (3.9). For $(\alpha, \beta) \in \mathbb{R}_+^2$, define

$$(3.11) \quad \hat{R} = \hat{R}(\alpha, \beta) = \sup\{r > 0 : u_1(x) > 0, u_2(x) > 0, x \in (0, r)\}.$$

Then $R(\alpha, \beta) \leq \hat{R}(\alpha, \beta)$ for any $(\alpha, \beta) \in \mathbb{R}_+^2$. Indeed it is easy to see that if $(\alpha, \beta) \in \mathcal{B} \cup \mathcal{R}$, then $R(\alpha, \beta) = \hat{R}(\alpha, \beta)$, while if $(\alpha, \beta) \in \mathcal{G} \cup \mathcal{Y}$, then $R(\alpha, \beta) < \hat{R}(\alpha, \beta)$ as

the solution can be extended beyond $R(\alpha, \beta)$ with $u_i(x) > 0$. For \hat{R} , we define

$$(3.12) \quad \begin{aligned} \mathcal{U} &= \{(\alpha, \beta) \in \mathbb{R}_+^2 : \hat{R} < \infty, u_1 > 0, u_2 > 0, x \in (0, \hat{R}), u_1(\hat{R}) > 0, u_2(\hat{R}) = 0\}, \\ \mathcal{V} &= \{(\alpha, \beta) \in \mathbb{R}_+^2 : \hat{R} < \infty, u_1 > 0, u_2 > 0, x \in (0, \hat{R}), u_1(\hat{R}) = 0, u_2(\hat{R}) > 0\}. \end{aligned}$$

Note that \mathcal{U} and \mathcal{V} are not restricted to I . Next we use this definition to show the behavior of solutions with initial values in III and IV .

Proposition 3.2. Let \mathcal{U} and \mathcal{V} be defined as in (3.12). Then \mathcal{U} and \mathcal{V} are open subsets of \mathbb{R}_+^2 such that $\mathcal{U} \supset \overline{III} \setminus \{(u_1^*, u_2^*)\}$ and a portion of I and II adjacent to III , and $\mathcal{V} \supset \overline{IV} \setminus \{(u_1^*, u_2^*)\}$ and a portion of I and II adjacent to IV .

Proof. We only prove the result for \mathcal{U} , and the one for \mathcal{V} also follows by symmetry. If $(\alpha, \beta) \in III$, then $f(\alpha, \beta) < 0$ and $g(\alpha, \beta) > 0$, and $u_1' > 0$ and $u_2' < 0$ for $x \in (0, \delta)$ from the equations. Let

$$(3.13) \quad R_1 = \sup\{r > 0 : u_1 > 0, u_2 > 0, u_1' > 0, u_2' < 0, x \in (0, r)\}.$$

If $R_1 < \infty$, we claim that $u_2(R_1) = 0$ and $u_1(R_1) > \alpha$. Indeed for $x \in (0, R_1)$, $u_1' > 0$ and $u_2' < 0$ hence $(u_1(x), u_2(x))$ remains in III so $u_1(x) > \alpha$ for $x \in (0, R_1]$. Also for $x \in (0, R_1]$,

$$(3.14) \quad u_1'(x) = - \int_0^x f(u_1, u_2) dt > 0, \quad \text{and} \quad u_2'(x) = - \int_0^x g(u_1, u_2) dt < 0.$$

Thus $u_2(R_1) = 0$. If $R_1 = \infty$, then $\lim_{x \rightarrow \infty} u_2(x) \geq 0$ exists and $u_2'(x) \rightarrow 0$ as $x \rightarrow \infty$, but $g(u_1(x), u_2(x)) > g(\alpha, \beta) \geq \delta_1 > 0$. From the second equation in (3.14), $u_2'(x) < -\delta_1 x$, that is a contradiction. Hence $R_1 = \infty$ is not possible, and $\hat{R} = R_1 < \infty$ satisfying $u_1(\hat{R}) > 0$ and $u_2(\hat{R}) = 0$. Thus $\mathcal{U} \supset III$.

If (α, β) is a boundary point of III but not (u_1^*, u_2^*) , without loss of generality, we assume that $f(\alpha, \beta) = 0$ and $g(\alpha, \beta) > 0$, then $u_2' < 0$ in $(0, \delta)$ and the orbit $(u_1(x), u_2(x))$ is in the interior of III for $x \in (0, \delta)$. Thus we can proceed the proof as above. The openness of \mathcal{U} follows from the continuous dependence of solutions of (3.6) on the initial conditions. Since $\mathcal{U} \supset \overline{III} \setminus \{(u_1^*, u_2^*)\}$, it also contains a portion in I and II adjacent to $\partial III \setminus \{(u_1^*, u_2^*)\}$ from the openness of \mathcal{U} . \square

The result in Proposition 3.2 shows that \mathcal{U} and \mathcal{V} are not empty, and the following result shows that there is an order for the elements in \mathcal{U} , \mathcal{V} and \mathcal{S} where the cooperativeness of the system is a key.

Proposition 3.3.

- (1) Suppose that $(\alpha_0, \beta_0) \in \mathcal{S}$, then $(\alpha, \beta_0) \in \mathcal{V}$ for any $0 < \alpha < \alpha_0$, and $(\alpha_0, \beta) \in \mathcal{V}$ for any $\beta > \beta_0$;
- (2) Suppose that $(\alpha_0, \beta_0) \in \mathcal{S}$, then $(\alpha, \beta_0) \in \mathcal{U}$ for any $\alpha > \alpha_0$, and $(\alpha_0, \beta) \in \mathcal{U}$ for any $0 < \beta < \beta_0$.

Proof. Again we only prove the first case. Assume that $(\alpha_0, \beta_0) \in \mathcal{S}$, then for $u_1(x) = u_1(x; \alpha_0, \beta_0)$, $u_2(x) = u_2(x; \alpha_0, \beta_0)$, we have $u_1, u_2 > 0$ and $u_1', u_2' < 0$ for $x \in (0, \hat{R}(\alpha_0, \beta_0))$. We claim that $\hat{R}(\alpha, \beta_0) < \hat{R}(\alpha_0, \beta_0)$. Define $\phi(x) =$

$u_1(x; \alpha_0, \beta_0) - u_1(x; \alpha, \beta_0)$ and $\psi(x) = u_2(x; \alpha_0, \beta_0) - u_2(x; \alpha, \beta_0)$, then (ϕ, ψ) satisfies

$$\begin{cases} \phi'' - \phi + h'_2(U_2)\psi = 0, & x \in (0, R_*), \\ \psi'' + h'_1(U_1)\phi - \psi = 0, & x \in (0, R_*), \\ \phi(0) = \alpha_0 - \alpha > 0, \quad \phi'(0) = 0, \\ \psi(0) = 0, \quad \psi'(0) = 0, \end{cases}$$

where $R_* = \min\{\hat{R}(\alpha, \beta_0), \hat{R}(\alpha_0, \beta_0)\}$, $U_1 = t_1(x)u_1(x; \alpha_0, \beta_0) + (1-t_1(x))u_1(x; \alpha, \beta_0) > 0$, and $U_2 = t_2(x)u_2(x; \alpha_0, \beta_0) + (1-t_2(x))u_2(x; \alpha, \beta_0) > 0$. Then by comparing with $(u'_1(x; \alpha_0, \beta_0), u'_2(x; \alpha_0, \beta_0))$ and using the same proof as in Lemma 2.2, we can prove that $\psi(x) < 0$ for $x \in (0, R_*]$, and ϕ has at most one zero in $(0, R_*]$. In fact, $\phi(x) > 0$ for $x \in (0, R_*]$, since the solution ϕ_3 of $\phi''_3 - \phi_3 = 0$ satisfying $\phi_3(0) = 1$ and $\phi'_3(0) = 0$ is clearly positive in $(0, R_*)$. This implies that at R_* , $u_1(R_*; \alpha_0, \beta_0) > u_1(R_*; \alpha, \beta_0)$ and $u_2(R_*; \alpha_0, \beta_0) < u_2(R_*; \alpha, \beta_0)$. Thus $R_* = \hat{R}(\alpha, \beta_0)$, $u_1(R_*; \alpha, \beta_0) = 0$, and $(\alpha, \beta_0) \in \mathcal{V}$. Similarly by comparing the solutions initiating from (α, β) and (α, β_0) , where $\alpha < \alpha_0$ and $\beta > \beta_0$, we can conclude that $(\alpha, \beta) \in \mathcal{V}$ as well. \square

The result in Proposition 3.3 suggests that the set \mathcal{S} of initial values generating the crossing solutions is on the boundary between the sets \mathcal{U} and \mathcal{V} . We now prove that is the case.

Proposition 3.4. For $\alpha > 0$, define

$$(3.15) \quad \Phi_1(\alpha) = \inf\{\beta > 0 : (\alpha, \beta) \in \mathcal{V}\}, \quad \Phi_2(\alpha) = \sup\{\beta > 0 : (\alpha, \beta) \in \mathcal{U}\}.$$

Then Φ_i ($i = 1, 2$) are well-defined. Moreover define

$$(3.16) \quad \alpha_* = \sup\{\alpha > 0 : H(\alpha, \Phi_1(\alpha)) \leq 0\},$$

then for $\alpha > \alpha_*$, $\Phi_1(\alpha) = \Phi_2(\alpha) \equiv \Phi(\alpha)$, where $\Phi : (\alpha_*, \infty) \rightarrow \mathbb{R}_+$ is a continuously differentiable, strictly increasing function. Moreover $\mathcal{S} = \{(\alpha, \Phi(\alpha)) : \alpha > \alpha_*\}$, $\mathcal{V} \supset \{(\alpha, \beta) : \alpha > \alpha_*, \beta > \Phi(\alpha)\}$, and $\mathcal{U} \supset \{(\alpha, \beta) : \alpha > \alpha_*, 0 < \beta < \Phi(\alpha)\}$.

Proof. First $\Phi_1(\alpha)$ is well-defined since for fixed $\alpha > 0$, $(\alpha, \beta) \in IV \subset \mathcal{V}$ if $\beta > 0$ is large enough; secondly $\Phi_1(\alpha) > 0$ for any $\alpha > 0$ since for fixed $\alpha > 0$, if $0 < \beta < \beta_0$, then $(\alpha, \beta) \in III \subset \mathcal{U}$ where $f(\alpha, \beta_0) = 0$. Similarly $\Phi_2(\alpha)$ is also well-defined.

From Proposition 3.3, $H(\alpha, \Phi_1(\alpha)) < 0$ for $\alpha > 0$ is small. And it is clear that $H(\alpha, \Phi_1(\alpha)) > 0$ when $\alpha > 0$ is large. Thus $\alpha_* > 0$ exists, and it is necessary that $H(\alpha_*, \Phi_1(\alpha_*)) = 0$. For $\alpha > \alpha_*$, $H(\alpha, \Phi_1(\alpha)) > 0$. We claim that if $\alpha > \alpha_*$, then $(\alpha, \Phi_1(\alpha)) \in \mathcal{S}$. To prove the claim, we fix $\alpha > \alpha_*$. Consider $\hat{R}(\alpha, \Phi_1(\alpha))$, there are two cases: (1) $\hat{R}(\alpha, \Phi_1(\alpha)) = \infty$, or (2) $\hat{R}(\alpha, \Phi_1(\alpha)) < \infty$. For Case (1), $\hat{R}(\alpha, \Phi_1(\alpha)) = \infty$, then $u_1(x), u_2(x) > 0$ for all $x \in [0, \infty)$, and there are two subcases: (1a) $R(\alpha, \Phi_1(\alpha)) = \infty$, or (1b) $R(\alpha, \Phi_1(\alpha)) < \infty$. For Case (1a), $(\alpha, \Phi_1(\alpha)) \in \mathcal{Q}$, but on the other hand $H(\alpha, \Phi_1(\alpha)) > 0$, which is a contradiction by Lemma 3.1. For Case (1b), let $R_2 = R(\alpha, \Phi_1(\alpha))$, then $u_1(R_2) = u_1^* > 0$ and $u_2(R_2) = u_2^* > 0$, and at least one of $u'_1(R_2)$ or $u'_2(R_2)$ is zero. If $u'_1(R_2) = u'_2(R_2) = 0$, then $f(u_1^*, u_2^*) = -u''_1(R_2) \leq 0$ and $g(u_1^*, u_2^*) = -u''_2(R_2) \leq 0$ so $(u_1^*, u_2^*) \in \bar{II}$ and $0 \geq H(u_1^*, u_2^*) = H_0(R_2) = H_0(0) = H(\alpha, \Phi_1(\alpha))$, that is a contraction with $H(\alpha, \Phi_1(\alpha)) > 0$. If only one of $u'_1(R_2)$ or $u'_2(R_2)$ is zero, without loss of generality, we assume that $u'_1(R_2) = 0$ and $u'_2(R_2) < 0$. Then we still have $f(u_1^*, u_2^*) = -u''_1(R_2) \leq 0$, and we must have $(u_1^*, u_2^*) \in III$. Then we can follow

the proof of Proposition 3.2 to show that $(u_1(x), u_2(x)) \in III$ for all $x > R_2$, which leads to $u_2(R_3) = 0$ for some $R_3 > R_2$, but that contradicts with $\hat{R}(\alpha, \Phi_1(\alpha)) = \infty$. This shows Case (1) cannot happen.

Hence we must have Case (2): $\hat{R}(\alpha, \Phi_1(\alpha)) < \infty$. Since $(\alpha, \Phi_1(\alpha))$ is a boundary point of V , from the continuity, $u_1(\hat{R}(\alpha, \Phi_1(\alpha))) = 0$, and $u_2(\hat{R}(\alpha, \Phi_1(\alpha))) \geq 0$. If $u_2(\hat{R}(\alpha, \Phi_1(\alpha))) > 0$, then $(\alpha, \Phi_1(\alpha)) \in V$, and V is open, so for some small $\epsilon > 0$, $(\alpha, \Phi_1(\alpha) - \epsilon) \in V$. That contradicts with the definition of Φ_1 . Thus we must have $u_2(\hat{R}(\alpha, \Phi_1(\alpha))) = 0$, and $(\alpha, \Phi_1(\alpha)) \in \mathcal{S}$.

Now since $(\alpha, \Phi_1(\alpha)) \in \mathcal{S}$ for $\alpha > \alpha_*$, then from Proposition 3.3, $(\alpha, \beta) \in \mathcal{V}$ for $\beta > \Phi_1(\alpha)$, and $(\alpha, \beta) \in \mathcal{U}$ for $0 < \beta < \Phi_1(\alpha)$. Similarly we can also prove that $(\alpha, \Phi_2(\alpha)) \in \mathcal{S}$ for $\alpha > \alpha_*$, which implies that we must have $\Phi_1(\alpha) = \Phi_2(\alpha)$ for $\alpha > \alpha_*$ from Proposition 3.3. And we denote it by $\Phi(\alpha)$ now. The continuity of $\Phi(\alpha)$ follows from the definition, and the differentiability of $\Phi(\alpha)$ follows from the differentiability of the solutions (u_1, u_2) on the initial values. Again from Proposition 3.3, $\Phi(\alpha)$ is strictly increasing in α since $(\alpha, \Phi_1(\alpha)) \in \mathcal{S}$.

We have proved that $\mathcal{S} \supset \{(\alpha, \Phi(\alpha)) : \alpha > \alpha_*\}$. We show that \mathcal{S} has no other elements other than the ones on $\{(\alpha, \Phi(\alpha)) : \alpha > \alpha_*\}$. For any $\alpha > \alpha_*$, there is no any elements of \mathcal{S} since $(\alpha, \beta) \in \mathcal{V}$ if $\beta > \Phi(\alpha)$, and $(\alpha, \beta) \in \mathcal{U}$ if $\beta < \Phi(\alpha)$. Same can be said for $\beta > \Phi(\alpha_*)$ from the symmetry. Hence possible elements of \mathcal{S} can only be in $W = (I \cap \{(\alpha, \beta) : 0 < \alpha \leq \alpha_*, 0 < \beta \leq \Phi(\alpha_*)\}) \setminus (\mathcal{U} \cup \mathcal{V})$. But for any $(\alpha, \beta) \in W$, $H(\alpha, \beta) \leq 0$. Thus $\mathcal{S} \cap W = \emptyset$ from Lemma 3.1. This proves $\mathcal{S} = \{(\alpha, \Phi(\alpha)) : \alpha > \alpha_*\}$. \square

Now we are at the position to classify all solutions of (3.2) and (3.3) with the aid of the general results established in previous sections:

Proposition 3.5. Let $\Phi(\alpha) : (\alpha_*, \infty) \rightarrow \mathbb{R}_+$ be defined as in Proposition 3.4. Define $R(\alpha)$ to be the common first zero of $u_1(x; \alpha, \Phi(\alpha))$ and $u_2(x; \alpha, \Phi(\alpha))$. Then $R(\alpha) : (\alpha_*, \infty) \rightarrow \mathbb{R}_+$ is a continuously differentiable, strictly decreasing function satisfying

$$(3.17) \quad \lim_{\alpha \rightarrow \alpha_*^+} R(\alpha) = \infty, \quad \text{and} \quad \lim_{\alpha \rightarrow \infty} R(\alpha) = 0,$$

and $\mathcal{Q} = (\alpha_*, \Phi(\alpha_*))$.

Proof. For $\alpha > \alpha_*$, we define the function $R(\alpha)$ by $R(\alpha) = R(\alpha, \Phi(\alpha)) = \hat{R}(\alpha, \Phi(\alpha))$, which is continuous and differentiable because of the continuous differentiability of $\Phi(\alpha)$ and the solution $u(x; \alpha, \beta)$ on initial values. We prove that $R(\alpha)$ is strictly decreasing. We differentiate $u_i(R(\alpha); \alpha, \Phi(\alpha)) = 0$ with respect to α for $i = 1, 2$, then

$$(3.18) \quad \begin{aligned} u_1'(R(\alpha))R'(\alpha) + \phi_1(R(\alpha)) + \Phi'(\alpha)\phi_2(R(\alpha)) &= 0, \\ u_2'(R(\alpha))R'(\alpha) + \psi_1(R(\alpha)) + \Phi'(\alpha)\psi_2(R(\alpha)) &= 0, \end{aligned}$$

where (ϕ_1, ψ_1) and (ϕ_2, ψ_2) are as defined in (2.6) and (2.7). Let $c = \Phi'(\alpha) > 0$, then (3.18) is equivalent to

$$(3.19) \quad u_1'(R(\alpha))R'(\alpha) = -A_c(R(\alpha)), \quad u_2'(R(\alpha))R'(\alpha) = -B_c(R(\alpha)),$$

where $(A_c, B_c) = (\phi_1, \psi_1) + c(\phi_2, \psi_2)$ is defined in (2.32). Since $u_1'(R(\alpha)) < 0$ and $u_2'(R(\alpha)) < 0$, then (3.19) implies that $A_c(R(\alpha))B_c(R(\alpha)) > 0$. Let c_1 and c_2 be defined as in (2.54). If $c = \Phi'(\alpha) > 0$ satisfies either $c \leq c_1$ or $c \geq c_2$, then $A_c(R(\alpha))B_c(R(\alpha)) \leq 0$ from Corollary 2.11. Hence $c = \Phi'(\alpha)$ satisfies $c_1 < c < c_2$.

Also from Corollary 2.11, we have $A_c(R(\alpha)) < 0$ and $B_c(R(\alpha)) < 0$, which implies $R'(\alpha) < 0$ by using (3.19). Since $R(\alpha)$ is decreasing, then $\lim_{\alpha \rightarrow \alpha_*^+} R(\alpha)$ exists. If the limit is finite, then $(\alpha_*, \Phi(\alpha_*)) \in \mathcal{S}$, but $H(\alpha_*, \Phi(\alpha_*)) = 0$, that is a contradiction with Lemma 3.1. Thus $\lim_{\alpha \rightarrow \alpha_*^+} R(\alpha) = \infty$. From the properties of h_i , $\Phi(\alpha) \rightarrow \infty$ as $\alpha \rightarrow \infty$. To prove $\lim_{\alpha \rightarrow \infty} R(\alpha) = 0$, it suffices to show that for any $R > 0$, there exists a positive solution for (3.2) with that $R > 0$. Indeed by using the existence result in [31] or [39], there exists a positive solution $u(x)$ for (3.2) with h_i satisfying (3.4) and (3.5).

Finally we prove that $\mathcal{Q} = (\alpha_*, \Phi(\alpha_*))$. We have shown that $\lim_{\alpha \rightarrow \alpha_*^+} R(\alpha) = \infty$, thus $(\alpha_*, \Phi(\alpha_*)) \in \mathcal{Q}$. By using similar proof as in that of Proposition 3.3, we can show that if $(\alpha_0, \beta_0) \in \mathcal{Q}$, then for any $(\alpha, \beta) \neq (\alpha_0, \beta_0)$, then $(\alpha, \beta) \in \mathcal{V}$ if $0 < \alpha \leq \alpha_0$ and $\beta \geq \beta_0$, and $(\alpha, \beta) \in \mathcal{U}$ if $\alpha \geq \alpha_0$ and $0 < \beta \leq \beta_0$. Suppose that there exists another $(\alpha^*, \beta^*) \in \mathcal{Q}$. Then $H(\alpha^*, \beta^*) = 0$, and $f(\alpha^*, \beta^*) > 0$, $g(\alpha^*, \beta^*) > 0$. Notice that $\partial H / \partial u_1 = g$ and $\partial H / \partial u_2 = f$, then either $\alpha^* > \alpha_*$, or $\beta^* > \Phi_1(\alpha_*)$. If $\alpha^* > \alpha_*$, then $\beta^* < \Phi(\alpha_*) < \Phi(\alpha^*)$, then $(\alpha^*, \beta^*) \in \mathcal{U}$ from the claim above. If $\beta^* > \Phi(\alpha_*)$, a similar contradiction can be reached. That is a contradiction. Thus \mathcal{Q} only has one element and $\mathcal{Q} = (\alpha_*, \Phi(\alpha_*))$. \square

We can now state our main result in this subsection about the existence and uniqueness of positive solutions to (3.2) and (3.3).

Theorem 3.6. *Suppose that h_i , ($i = 1, 2$), satisfy (3.4) and (3.5), then for any $R > 0$, (3.2) has a unique positive solution $(u_1(x; R), u_2(x; R))$. If $R_1 > R_2$, then $u_1(0; R_1) < u_1(0; R_2)$ and $u_2(0; R_1) < u_2(0; R_2)$. Moreover (3.3) has a unique solution (U_1, U_2) .*

Proof. For any $R > 0$, from Proposition 3.5, there exists a unique $\alpha \in (\alpha_*, \infty)$ such that $R(\alpha) = R$, hence the solution $(u_1(x; \alpha, \Phi(\alpha)), u_2(x; \alpha, \Phi(\alpha)))$ of (3.6) is a positive solution of (3.2). From Proposition 3.4, $\mathcal{S} = \{(\alpha, \Phi(\alpha)) : \alpha > \alpha_*\}$, which shows the uniqueness of positive with $R > 0$. From Proposition 3.5, the unique solution of (3.3) is the one of (3.6) with initial value $(\alpha_*, \Phi(\alpha_*))$. \square

We remark that as a limit of elements in \mathcal{S} , the unique ground state solution with $(\alpha, \beta) \in \mathcal{Q}$ is necessarily strictly decreasing, i.e. $u_1'(x) < 0$ and $u_2'(x) < 0$ for $x \in (0, \infty)$. For (α, β) satisfying $H(\alpha, \beta) \leq 0$, there are other positive solutions of (3.6) defined for all $x > 0$, but they are oscillatory around (u_1^*, u_2^*) and not converge to $(0, 0)$. It is possible to classify all these solutions using similar method here, but we will not pursue that here.

Example 3.7. Consider

$$(3.20) \quad \begin{cases} u_1'' - u_1 + u_2^q = 0, & -R < x < R, \\ u_2'' - u_2 + u_1^p = 0, & -R < x < R, \\ u_1(x) > 0, \quad u_2(x) > 0, & -R < x < R, \\ u_1(\pm R) = u_2(\pm R) = 0, & \end{cases}$$

where $p, q > 1$. Then the conditions (3.4) and (3.5) are satisfied. Our result shows that for any $R > 0$, (3.20) has a unique solution, which is even and decreasing in $(0, R)$, and there is a unique (up to a translation) ground state solution of (3.20) which is defined on \mathbb{R} . Equations like (3.20) was considered in [31, 33, 39], in which




FIGURE 1. Numerical bifurcation diagram for (3.20) for $q = 2$ and $p = 3$. Here the horizontal axis is $u_1(0) = \alpha \in [0, 5]$, the vertical axis is $u_2(0) = \beta \in [0, 5]$. Regions *II*, *III*, and *IV* are colored in cyan, while region *I* are according to the behavior of solution at $R(\alpha, \beta)$: \mathcal{B} (blue), \mathcal{G} (green), \mathcal{R} (red), \mathcal{Y} (yellow). The curves $f(u_1, u_2) = 0$ (bordering \mathcal{G}), $g(u_1, u_2) = 0$ (bordering \mathcal{Y}), and $H(u_1, u_2) = 0$ are plotted in black.

the existence and symmetry of the positive solution for a general bounded domain Ω were proved. Our result appears to the first uniqueness result for such superlinear Hamiltonian elliptic systems. The existence and uniqueness of positive solution to (3.20) for the case of $pq < 1$ and Ω being a general bounded domain was proved in [13]. The uniqueness of the positive solution to (3.20) for $pq \geq 1$ and $p > 1 > q$ (or $q > 1 > p$) is still not known.

A numerical bifurcation diagram of (3.20) is shown in Figure 1. One can see that $U \supset III \cup \mathcal{G} \cup \mathcal{R}$, and $V \supset IV \cup \mathcal{Y} \cup \mathcal{B}$. The boundary between \mathcal{B} and \mathcal{R} is the set \mathcal{S} of crossing solutions, and the unique common point of \mathcal{G} , \mathcal{R} , \mathcal{Y} , \mathcal{B} and also $H = 0$ is the element in \mathcal{Q} . The numerical bifurcation diagram is plotted using method described in [36] and `Matlab`.

Remark 3.8.

- (1) Our uniqueness result for (3.20) with $p, q > 1$ can also be viewed as a generalization (in the case of $n = 1$) of the uniqueness of positive solution to a well-known scalar equation:

$$(3.21) \quad \begin{cases} \Delta u - u + u^p = 0, & x \in \mathbb{R}^n, \\ u(x) > 0, & x \in \mathbb{R}^n, \\ u(x) \rightarrow 0, & |x| \rightarrow \infty, \end{cases}$$

where $1 < p < \frac{n+2}{n-2}$ for $n \geq 3$ and $1 < p < \infty$ for $n = 1, 2$. The solution of (3.21) serves as the basic building block of spike layer solutions in a singularly perturbed elliptic problem arising from the pattern formation of hydra (see [46]). The uniqueness of the solution of (3.21) was first proved by

Coffman [22] when $n = 3$, and by Kwong [44] for general subcritical power nonlinearity. Simplifications of proof and generalizations can be found in, for example, [5, 10, 45, 52, 53, 54, 62].

- (2) Korman [42] proved the uniqueness of positive solution for a Hamiltonian system

$$(3.22) \quad \begin{cases} u'' + \lambda f(v) = 0, & -1 < x < 1 \\ v'' + \lambda g(u) = 0, & -1 < x < 1, \\ u(\pm 1) = v(\pm 1) = 0. \end{cases}$$

under the conditions that $f(t) > 0$, $g(t) > 0$, $f'(t) > 0$, $g'(t) > 0$ and $tf'(t) \geq f(t)$ and $tg'(t) \geq g(t)$ for all $t > 0$. Note that these $f(t)$ and $g(t)$ are superlinear, but they are positive thus no ground state is possible here. Our approach in this subsection can be adapted to give a proof of result in [42].

3.2. A gradient quadratic Schrödinger system. In this subsection we consider the uniqueness of positive solutions of the $\chi^{(2)}$ Second-Harmonic Generation (SHG) equations from the nonlinear optics models (see [48, 69, 70]).

$$(3.23) \quad \begin{cases} u_1'' - bu_1 + u_1u_2 = 0, & x \in (0, R), \\ u_2'' + u_1^2/2 - cu_2 = 0, & x \in (0, R), \\ u_1'(0) = u_2'(0) = u_1(R) = u_2(R) = 0, \end{cases}$$

where $b, c > 0$. The system (3.23) also appears in the study of gravity water waves, and it has been studied from the mathematical point of view in the work [23, 24, 34, 35]. When $c = 0$ in (3.23), it is also called Schrödinger-Newton equation (see [55, 66]). The corresponding equation of (3.23) for $x \in (0, \infty)$ is

$$(3.24) \quad \begin{cases} u_1'' - bu_1 + u_1u_2 = 0, & x \in (0, \infty), \\ u_2'' + u_1^2/2 - cu_2 = 0, & x \in (0, \infty), \\ u_1(x) > 0, u_2(x) > 0, u_1'(x) < 0, u_2'(x) < 0, & x \in (0, \infty), \\ u_1'(0) = u_2'(0), \end{cases}$$

Here we consider the initial value problem (2.3) with

$$(3.25) \quad f(u_1, u_2) = -bu_1 + u_1u_2, \quad g(u_1, u_2) = \frac{1}{2}u_1^2 - cu_2.$$

It is easy to verify that (f, g) is superlinear but not strongly superlinear, and it is also weakly sublinear. Clearly it is also cooperative. Define

$$(3.26) \quad F(u_1, u_2) = -\frac{b}{2}u_1^2 - \frac{c}{2}u_2^2 + \frac{1}{2}u_1^2u_2.$$

Then we can see that (3.23) is a gradient system with energy function defined in (1.16) and (3.26).

We first consider the case $c > 0$ for (3.23). For the f and g defined as in (3.25), we use the definitions of the set I, II, III, IV as in (3.8), and $\mathcal{U}, \mathcal{V}, \mathcal{S}, \mathcal{Q}$ as in (3.9) and (3.12). Then the results in Lemma 3.1 hold except $H(\alpha, \beta)$ is replaced by $F(\alpha, \beta)$. Next the results in Propositions 3.2, 3.3 and 3.4 also hold with essentially the same proof, and this leads to the result in Proposition 3.5 in the exact same wording except $H(\alpha, \beta)$ is replaced by $F(\alpha, \beta)$. Note that here (f, g) is again superlinear and weakly sublinear as in the proof of Proposition 3.5. The existence of a positive

solution of (3.23) with $c > 0$ for any $R > 0$ can be proved via a standard variational method using the Mountain-pass lemma (see [2]) for the functional

$$E(u_1, u_2) = \int_{-R}^R \left(\frac{1}{2}[u_1'(x)]^2 + \frac{1}{2}[u_2'(x)]^2 - F(u_1(x), u_2(x)) \right) dx,$$

where F is defined by (3.26), and $(u_1, u_2) \in [H_0^1(-R, R)]^2$. Indeed one can show that (i) Palais-Smale condition is satisfied for E ; (ii) (u_1, u_2) is a local minimum of the functional E , and (iii) $E(t\phi_1, t\phi_1) < 0$ for $t > 0$ large and $\phi_1(x) = \cos(\pi x/(2R)) > 0$. Hence the Mountain-pass lemma can be applied. The existence of positive solution defined for $x \in \mathbb{R}$ was proved in [69] by using again the variational approach but with the concentration-compactness principle due to the lack of compactness. Again the element in \mathcal{Q} is unique as the set $F(\alpha, \beta) = 0$ in the region I is a curve $\{(\alpha, \mathcal{F}(\alpha))\}$ such that $\mathcal{F}'(\alpha) < 0$. In summary we obtain the following uniqueness result for the solutions of (3.23):

Theorem 3.9. *Suppose that $b, c > 0$, then for any $R > 0$, (3.23) has a unique positive solution $(u_1(x; R), u_2(x; R))$. If $R_1 > R_2$, then $u_1(0; R_1) < u_1(0; R_2)$ and $u_2(0; R_1) < u_2(0; R_2)$. Moreover (3.24) has a unique solution (U_1, U_2) .*




FIGURE 2. Numerical bifurcation diagram for (3.23) for $b = 1$ and $c = 0.8$. Here the horizontal axis is $u_1(0) = \alpha \in [0, 5]$, the vertical axis is $u_2(0) = \beta \in [0, 5]$. Regions II , III , and IV are colored in cyan, while region I are according to the behavior of solution at $R(\alpha, \beta)$: \mathcal{B} (blue), \mathcal{G} (green), \mathcal{R} (red), \mathcal{Y} (yellow). The curves $f(u_1, u_2) = 0$ (bordering \mathcal{G}), $g(u_1, u_2) = 0$ (bordering \mathcal{Y}), and $F(u_1, u_2) = 0$ are plotted in black.

Remark 3.10. The uniqueness of the positive solution of (3.24) has been proved in Lopes [49] by using his earlier result [48] that the linearized problem of (3.24) has zero as a simple eigenvalue and has exactly one negative eigenvalue when $b, c > 0$. Here we obtain not only the uniqueness of ground state solution but also the uniqueness of crossing solution on the bounded intervals. Also our approach is more general and applicable to more problems.

Next we consider the case $c = 0$ for (3.23). Then (3.25) is written as

$$(3.27) \quad f(u_1, u_2) = -bu_1 + u_1u_2, \quad g(u_1, u_2) = \frac{1}{2}u_1^2.$$

We still use the notations defined in (3.8) and (3.9), but we notice that here $g > 0$ for all $(u_1, u_2) \in \mathbb{R}_+^2$, so $II = IV = \emptyset$. Again for a solution of (2.3) with $(\alpha, \beta) \in I$, we define R to be the right endpoint of the maximal interval $(0, R(\alpha, \beta))$ so that $u_i(x) > 0$ and $u_i'(x) < 0$ in $(0, R(\alpha, \beta))$, and we define \hat{R} to be the right endpoint of the maximal interval $(0, \hat{R}(\alpha, \beta))$ so that $u_i(x) > 0$ in $(0, \hat{R}(\alpha, \beta))$, $i = 1, 2$. The following lemma characterizes the sets \mathcal{U} , \mathcal{V} , \mathcal{S} and \mathcal{Q} for the case of (3.23) with $c = 0$:

Lemma 3.11. *Suppose that f and g are defined as in (3.27), and the sets \mathcal{U} , \mathcal{V} , \mathcal{S} and \mathcal{Q} are defined as in (3.12) and (3.9).*

- (1) $\mathcal{V} \neq \emptyset$, and if $(\alpha, \beta) \in \mathcal{V}$, then $u_1'(x) < 0$, $u_2'(x) < 0$ for $x \in (0, \hat{R}(\alpha, \beta))$ (so $R(\alpha, \beta) = \hat{R}(\alpha, \beta)$).
- (2) $\mathcal{U} \neq \emptyset$, and if $(\alpha, \beta) \in \mathcal{U}$, then $u_2'(x) < 0$ for $x \in (0, \hat{R}(\alpha, \beta))$ and, either $u_1'(x) < 0$ for $x \in (0, \hat{R}(\alpha, \beta))$ or there exists $R_1 > 0$ such that $u_1'(x)(x - R_1) > 0$ for $x \in (0, \hat{R}(\alpha, \beta)) \setminus \{R_1\}$.
- (3) $\mathcal{Q} = \emptyset$ and $I = \mathcal{U} \cup \mathcal{V} \cup \mathcal{S}$.
- (4) If $(\alpha, \beta) \in \mathcal{S}$, then $u_1'(x) < 0$, $u_2'(x) < 0$ for $x \in (0, R(\alpha, \beta)]$.

Proof. 1. First we prove that if $(\alpha, \beta) \in \mathcal{V}$, then $u_1'(x) < 0$, $u_2'(x) < 0$ for $x \in (0, \hat{R})$ hence $R = \hat{R}$. Since $u_1(x) > 0$ for $x \in (0, \hat{R})$, then $u_2'(x) < 0$ for $x \in (0, \hat{R})$ from $u_2''(x) = -\frac{1}{2}u_1^2(x) < 0$. Since $\beta > b$, then $u_1'(x) < 0$ for x near 0. Suppose that there exists $R_2 \in (0, \hat{R})$ such that $u_1'(x) < 0$ for $x \in (0, R_2)$ and $u_1'(R_2) = 0$, then $x = R_2$ is a local minimum of $u_1(x)$, $u_1''(R_2) \geq 0$ and u_1 is increasing for $x \in (R_2, R_2 + \epsilon)$. Since $u_1(\hat{R}) = 0$, then there exists $R_3 \in (R_2, \hat{R})$ such that $u(x)$ achieves a local maximum at $x = R_3$. $u_1'(x) > 0$ for $x \in (R_2, R_3)$. This follows that $(u_2(R_2) - b)u_1(R_2) = -u_1''(R_2) \leq 0$ and $(u_2(R_3) - b)u_1(R_3) = -u_1''(R_3) \geq 0$, and one of the inequalities must be strict since $u_1(R_3) > u_1(R_2)$. Then $u_2(R_3) \geq b \geq u_2(R_2)$ and $u_2(R_3) > u_2(R_2)$, which contradicts with $u_2'(x) < 0$ for $x \in (0, \hat{R})$. Therefore $u_1'(x) < 0$ for $x \in (0, \hat{R})$.

Next we prove that $\mathcal{V} \neq \emptyset$. Indeed we prove that any fixed $\beta > b$, there exists a constant $\delta_1 = \delta_1(\beta) > 0$ such that for $0 < \alpha < \delta_1$, $(\alpha, \beta) \in \mathcal{V}$. Let $\beta - b = 2\epsilon$ and let $v_1(x)$ be the solution to

$$(3.28) \quad \begin{cases} v''(x) + \epsilon v(x) = 0, & x > 0, \\ v(0) = \alpha, v'(0) = 0. \end{cases}$$

Since Eq. (3.28) is a linear equation, then there exist $R_4 = \pi/(2\sqrt{\epsilon}) > 0$ such that $v_1(R_4) = 0$ and $v_1(x) > 0$, $v_1'(x) < 0$ for $x \in (0, R_4)$. Let $(u_1(x), u_2(x))$ be a solution of (2.3) with initial value (α, β) satisfying $0 < \alpha < 2\sqrt{\epsilon}/R_4 = 4\epsilon^2/\pi$. We define $R_5 = \sup\{x > 0 : u_1(s) > 0, u_2(s) > b + \epsilon, \text{ for all } s \in (0, x)\}$, and $R_6 = \min\{R_5, R_4\}$. Then from the second equation of (2.3) and $u_1'(x) < 0$ in $(0, R_5)$, we have for any $x \in (0, R_6)$,

$$(3.29) \quad -u_2'(x) = \int_0^x \frac{1}{2}u_1^2(s) ds \leq \frac{1}{2} \int_0^x \alpha^2 ds = \frac{1}{2}\alpha^2 x.$$

Integrating (3.29) again on $[0, R_6]$, we obtain

$$-u_2(R_6) + \beta \leq \frac{\alpha^2 R_6^2}{4} < \varepsilon,$$

from the assumption on α . This implies that $u_2(R_6) > \beta - \varepsilon = b + \varepsilon$. On the other hand, for $r \in (0, R_6)$,

$$u_1''(x) = -(u_2(x) - b)u_1(x) < -\varepsilon u_1(x).$$

By the comparison principle, $u_1(x) \leq v_1(x)$ for all $x > 0$. Since $v_1(R_4) = 0$, then $u_1(x)$ must reach zero for some $x < R_4$. Therefore $R_6 = R_5 < R_4$ and $u_1(R_5) = 0$ and $u_2(x) > b + \varepsilon$ for $x \in (0, R_5)$. This prove that $(\alpha, \beta) \in \mathcal{V}$ when $0 < \alpha < 4\varepsilon^2/\pi$.

2. First we prove the monotonicity of $u_1(x)$ and $u_2(x)$ when $(\alpha, \beta) \in \mathcal{U}$. Again since $u_1(x) > 0$ for $x \in (0, \hat{R})$, then $u_2'(x) < 0$ for $x \in (0, \hat{R})$ from $u_2''(x) = -\frac{1}{2}u_1^2(x) < 0$. Since $\beta > b$, then $u_1'(x) < 0$ for x near 0. By the arguments above, for $x \in (0, \hat{R})$, one cannot have $0 < R_2 < R_3 < \hat{R}$ such that $x = R_2$ is a local minimum and $x = R_3$ is a local maximum of $u_1(x)$. Hence either $u_1'(x) < 0$ for $x \in (0, \hat{R})$ or there exists $R_1 > 0$ such that $u_1'(x)(x - R_1) > 0$ for $x \in (0, \hat{R}) \setminus \{R_1\}$.

Next we prove that $\mathcal{U} \neq \emptyset$. Indeed we prove that any fixed $\alpha > 0$, there exists a constant $\delta_2 = \delta_2(\alpha) > 0$ such that for $b < \beta < \delta_2$, $(\alpha, \beta) \in \mathcal{U}$. For a fixed $\alpha > 0$, if $\beta = b$, then $u_1'(x) > 0$ and $u_2'(x) < 0$ for small $x > 0$. For a small $\varepsilon > 0$, let $R_7 > 0$ such that $u_1(R_7; \alpha, b) = \alpha + 2\varepsilon$. Then from the continuous dependence of (2.3) on the initial value, there exists $\delta_2 > 0$ such that when $b < \beta < b + \delta_2$, $u_1(R_7; \alpha, \beta) > \alpha + \varepsilon$. In particular this implies that there exists $R_1 \in (0, R_7)$ such that $u_1'(x; \alpha, \beta)(x - R_1) > 0$ for $x \in (0, \hat{R}) \setminus \{R_1\}$. Then for this $u_1(x)$, we have $u_1(x) \geq u_1(R_1)$ for $x \in (0, \hat{R})$, which implies that

$$(3.30) \quad -u_2'(x) = \int_0^x \frac{1}{2} u_1^2(s) ds \geq \frac{1}{2} \int_0^x u_1^2(R_1) ds = \frac{1}{2} u_1^2(R_1)x.$$

By integrating (3.30) on $[0, r]$ for $r > 0$, we obtain $u_2(r) \leq \beta - u_1^2(R_1)r^2/4$. Then for $r > 2\sqrt{\beta}/u_1(R_1)$, we have $u_2(r) \leq 0$ which implies that $(\alpha, \beta) \in \mathcal{U}$.

3. Suppose that $(\alpha, \beta) \in \mathcal{Q}$, then $u_2(x)$ is a positive subharmonic function in \mathbb{R} by an even extension. Hence $u_2(x)$ must be a constant and $u_1(x) \equiv 0$ which contradicts with $\alpha > 0$. Therefore $\mathcal{Q} = \emptyset$. Since for any $(\alpha, \beta) \in I$, $u_1(x)$ is either strictly decreasing in $(0, \hat{R})$, or there exists $R_1 > 0$ such that $u_1'(x)(x - R_1) > 0$ for $x \in (0, \hat{R}) \setminus \{R_1\}$, then $u_2(x)$ is always decreasing in $(0, \hat{R})$. Also we have proved that $\mathcal{Q} = \emptyset$. Then \hat{R} must be finite, and either $u_1(\hat{R}) = 0$ or $u_2(\hat{R}) = 0$. This proves $I = \mathcal{U} \cup \mathcal{V} \cup \mathcal{S}$, and also when $(\alpha, \beta) \in \mathcal{S}$, we must have $u_1'(x) < 0$, $u_2'(x) < 0$ for $x \in (0, \hat{R}]$. \square

Secondly we consider the linearized equations (2.6) and (2.7). Here we have

$$(3.31) \quad f_1 = -b + u_2(x), \quad f_2 = u_1(x), \quad g_1 = u_1(x), \quad g_2 = 0.$$

It is clear that the system is cooperative, and (f, g) is superlinear and weakly sublinear. Then the solutions (ϕ_i, ψ_i) ($i = 1, 2$) of the linearized equation are of one sign. That is

$$(3.32) \quad \phi_1(x) > 0, \quad \psi_1(x) < 0, \quad \phi_2(x) < 0, \quad \psi_2(x) > 0.$$

Similar to the proof in subsection 3.1, we are able to show the characterization of the set \mathcal{U} , \mathcal{V} and \mathcal{S} as in Proposition 3.3 also hold here. Since the statement is exactly the same, we omit it here.

Now we can classify all solutions of the initial value problem (2.3) with (f, g) defined in (3.27).

Proposition 3.12. Consider equation (2.3) with $c = 0$, and (f, g) is defined in (3.27).

- (1) There exists a strictly increasing function $\Phi : (0, \infty) \rightarrow (b, \infty)$ satisfying

$$(3.33) \quad \lim_{\alpha \rightarrow 0^+} \Phi(\alpha) = b \quad \text{and} \quad \lim_{\alpha \rightarrow \infty} \Phi(\alpha) = \infty,$$

such that

$$(3.34) \quad \begin{aligned} \mathcal{S} &= \{(\alpha, \Phi(\alpha)) : \alpha > 0\}, \\ \mathcal{V} &= \{(\alpha, \beta) : \alpha > 0, \beta > \Phi(\alpha)\}, \\ \mathcal{U} &= \{(\alpha, \beta) : \alpha > 0, b < \beta < \Phi(\alpha)\}. \end{aligned}$$

- (2) Define $R(\alpha)$ to be the common first zero of $u_1(x; \alpha, \Phi(\alpha))$ and $u_2(x; \alpha, \Phi(\alpha))$. Then $R(\alpha) : (0, \infty) \rightarrow (0, \infty)$ is a continuously differentiable, strictly decreasing function satisfying $\lim_{\alpha \rightarrow 0^+} R(\alpha) = \infty$ and $\lim_{\alpha \rightarrow \infty} R(\alpha) = 0$.

Proof. 1. Similar to the proof of Proposition 3.4, for $\alpha > 0$, we define Φ_1 and Φ_2 as in (3.15). Again from the continuous dependence of solutions of (2.3) on the initial conditions, both of \mathcal{U} and \mathcal{V} are open subsets of I . From Lemma 3.11, $\mathcal{Q} = \emptyset$ and for $\alpha > 0$, we can show that $(\alpha, \Phi_i(\alpha)) \in \mathcal{S}$ for $i = 1, 2$ using the same argument as in the proof of Proposition 3.4, therefore $\Phi_1(\alpha) = \Phi_2(\alpha)$ (which we define as $\Phi(\alpha)$). This proves the characterization in (3.34). The properties in Proposition 3.3 imply that $\Phi(\alpha)$ must be strictly increasing. From the proof in Lemma 3.11, the limits in (3.33) hold.

2. The proof for the properties of $R(\alpha)$ is similar to that in the proof of Proposition 3.5. \square

The existence of positive solution of (3.23) with $c = 0$ and any $R > 0$ can be proved using the same variational method as the $c > 0$ case. Hence we have the following result:

Theorem 3.13. *Suppose that $b > 0$ and $c = 0$, then for any $R > 0$, (3.23) has a unique positive solution $(u_1(x; R), u_2(x; R))$. If $R_1 > R_2$, then $u_1(0; R_1) < u_1(0; R_2)$ and $u_2(0; R_1) < u_2(0; R_2)$. Moreover (3.24) has no positive solution.*

Acknowledgement. The authors thank Professor Chang-Shou Lin for his helpful discussions regarding this project. The authors also thank two anonymous reviewers for their helpful comments which improve the manuscript. This work was done during several visits by J.-P. Shi to National Center for Theoretical Sciences (NCTS) in National Tsing-Hua University and National Central University (2007, 2010, 2013, 2014), visit of Y.-L. An (2014) and J.-L. Chern (2008, 2012) to College of William and Mary. They thank these institutes for their hospitality and support.

REFERENCES

1. A. Ambrosetti and E. Colorado, *Standing waves of some coupled nonlinear Schrödinger equations*, J. Lond. Math. Soc. (2) **75** (2007), no. 1, 67–82. MR 2302730 (2008f:35369)
2. A. Ambrosetti and P. H. Rabinowitz, *Dual variational methods in critical point theory and applications*, J. Functional Analysis **14** (1973), 349–381. MR 0370183
3. W. O. Amrein, A. M. Hinz, and D. B. Pearson (eds.), *Sturm-Liouville theory: Past and present*, Birkhäuser Verlag, Basel, 2005. MR 2132131 (2006a:34003)




FIGURE 3. Numerical bifurcation diagram for (3.23) for $b = 1$ and $c = 0$. Here the horizontal axis is $u_1(0) = \alpha \in [0, 5]$, the vertical axis is $u_2(0) = \beta \in [0, 5]$. Regions *II*, *III*, and *IV* are colored in cyan, while region *I* are according to the behavior of solution at $R(\alpha, \beta)$: \mathcal{B} (blue), \mathcal{G} (green), \mathcal{R} (red). The curve $f(u_1, u_2) = 0$ (bordering \mathcal{G}) is plotted in black.

4. Y.-L. An, *Uniqueness of positive solutions for a class of elliptic systems*, J. Math. Anal. Appl. **322** (2006), no. 2, 1071–1082. MR 2250636 (2007c:35044)
5. P. W. Bates and J.-P. Shi, *Existence and instability of spike layer solutions to singular perturbation problems*, J. Funct. Anal. **196** (2002), no. 2, 211–264. MR 1943093 (2003k:35045)
6. H. Berestycki, L. Nirenberg, and S. R. S. Varadhan, *The principal eigenvalue and maximum principle for second-order elliptic operators in general domains*, Comm. Pure Appl. Math. **47** (1994), no. 1, 47–92. MR 1258192 (95h:35053)
7. I. Birindelli, E. Mitidieri, and G. Sweers, *Existence of the principal eigenvalue for cooperative elliptic systems in a general domain*, Differential Equations **35** (1999), no. 3, 326–334. MR 1726799 (2001c:35169)
8. J. Busca and B. Sirakov, *Symmetry results for semilinear elliptic systems in the whole space*, J. Differential Equations **163** (2000), no. 1, 41–56. MR 1755067 (2001m:35100)
9. H. Chan, C.-C. Fu, and C.-S. Lin, *Non-topological multi-vortex solutions to the self-dual Chern-Simons-Higgs equation*, Comm. Math. Phys. **231** (2002), no. 2, 189–221. MR 1946331 (2003m:58026)
10. C.-C. Chen and C.-S. Lin, *Uniqueness of the ground state solutions of $\Delta u + f(u) = 0$ in \mathbf{R}^n , $n \geq 3$* , Comm. Partial Differential Equations **16** (1991), no. 8-9, 1549–1572. MR 1132797 (92j:35048)
11. Z.-J. Chen and W.-M. Zou, *Positive least energy solutions and phase separation for coupled Schrödinger equations with critical exponent*, Arch. Ration. Mech. Anal. **205** (2012), no. 2, 515–551. MR 2947540
12. Z.-Y. Chen and J.-L. Chern, *Uniqueness of topological multivortex solutions in the Maxwell-Chern-Simons model*, J. Funct. Anal. **270** (2016), no. 6, 2073–2125. MR 3460235
13. Z.-Y. Chen, J.-L. Chern, J.-P. Shi, and Y.-L. Tang, *On the uniqueness and structure of solutions to a coupled elliptic system*, J. Differential Equations **249** (2010), no. 12, 3419–3442. MR 2737436 (2011m:35082)
14. Z.-Y. Chen, J.-L. Chern, and Y.-L. Tang, *On the solutions to a Liouville-type system involving singularity*, Calc. Var. Partial Differential Equations **43** (2012), no. 1-2, 57–81. MR 2860403 (2012j:35083)

15. J.-L. Chern, Z.-Y. Chen, J.-H. Chen, and Y.-L. Tang, *On the classification of standing wave solutions for the Schrödinger equation*, Comm. Partial Differential Equations **35** (2010), no. 2, 275–301. MR 2748625 (2011k:35069)
16. J.-L. Chern, Z.-Y. Chen, and C.-S. Lin, *Uniqueness of topological solutions and the structure of solutions for the Chern-Simons system with two Higgs particles*, Comm. Math. Phys. **296** (2010), no. 2, 323–351. MR 2608118 (2011i:58020)
17. J.-L. Chern, Z.-Y. Chen, Y.-L. Tang, and C.-S. Lin, *Uniqueness and structure of solutions to the Dirichlet problem for an elliptic system*, J. Differential Equations **246** (2009), no. 9, 3704–3714. MR 2515175 (2010f:35072)
18. J.-L. Chern, Y.-L. Tang, C.-S. Lin, and J.-P. Shi, *Existence, uniqueness and stability of positive solutions to sublinear elliptic systems*, Proc. Roy. Soc. Edinburgh Sect. A **141** (2011), no. 1, 45–64. MR 2773438 (2012f:35116)
19. J.-L. Chern and S.-G. Yang, *Evaluating solutions on an elliptic problem in a gravitational gauge field theory*, J. Funct. Anal. **265** (2013), no. 7, 1240–1263. MR 3073254
20. ———, *The non-topological fluxes of a two-particle system in the Chern-Simons theory*, J. Differential Equations **256** (2014), no. 10, 3417–3439. MR 3177901
21. ———, *Uniqueness of topological vortex in a self-dual Maxwell-Chern-Simons-Higgs system*, Indiana Univ. Math. J. **64** (2015), no. 6, 1747–1766. MR 3436234
22. C. V. Coffman, *Uniqueness of the ground state solution for $\delta u - u + u^3 = 0$ and a variational characterization of other solutions*, Arch. Ration. Mech. Anal. **46** (1972), 81–95.
23. E. Colorado, *Existence of bound and ground states for a system of coupled nonlinear Schrödinger-KdV equations*, C. R. Math. Acad. Sci. Paris **353** (2015), no. 6, 511–516. MR 3348984
24. ———, *On the existence of bound and ground states for some coupled nonlinear Schrödinger-Korteweg-de Vries equations*, Adv. Nonlinear Anal. DOI: [10.1515/anona-2015-0181](https://doi.org/10.1515/anona-2015-0181) (2016).
25. R.-H. Cui, P. Li, J.-P. Shi, and Y.-W. Wang, *Existence, uniqueness and stability of positive solutions for a class of semilinear elliptic systems*, Topol. Methods Nonlinear Anal. **42** (2013), no. 1, 91–104. MR 3155616
26. R.-H. Cui, Y.-W. Wang, and J.-P. Shi, *Uniqueness of the positive solution for a class of semilinear elliptic systems*, Nonlinear Anal. **67** (2007), no. 6, 1710–1714. MR 2326023 (2008g:35035)
27. R. Dalmasso, *Existence and uniqueness of positive radial solutions for the Lane-Emden system*, Nonlinear Anal. **57** (2004), no. 3, 341–348. MR 2064095 (2005b:35093)
28. D. G. de Figueiredo, *Monotonicity and symmetry of solutions of elliptic systems in general domains*, NoDEA Nonlinear Differential Equations Appl. **1** (1994), no. 2, 119–123. MR 1273345
29. ———, *Semilinear elliptic systems*, Nonlinear functional analysis and applications to differential equations (Trieste, 1997), World Sci. Publ., River Edge, NJ, 1998, pp. 122–152. MR 1703530 (2000e:35043)
30. ———, *Semilinear elliptic systems: existence, multiplicity, symmetry of solutions*, Handbook of differential equations: stationary partial differential equations. Vol. V, Handb. Differ. Equ., Elsevier/North-Holland, Amsterdam, 2008, pp. 1–48. MR 2497896 (2010j:35126)
31. D. G. de Figueiredo and P. L. Felmer, *On superquadratic elliptic systems*, Trans. Amer. Math. Soc. **343** (1994), no. 1, 99–116. MR 1214781 (94g:35072)
32. D. G. de Figueiredo and E. Mitidieri, *Maximum principles for linear elliptic systems*, Rend. Istit. Mat. Univ. Trieste **22** (1990), no. 1-2, 36–66 (1992). MR 1210477
33. D. G. De Figueiredo and J.-F. Yang, *Decay, symmetry and existence of solutions of semilinear elliptic systems*, Nonlinear Anal. **33** (1998), no. 3, 211–234. MR 1617988 (99d:35047)
34. J.-P. Dias, M. Figueira, and F. Oliveira, *Existence of bound states for the coupled Schrödinger-KdV system with cubic nonlinearity*, C. R. Math. Acad. Sci. Paris **348** (2010), no. 19-20, 1079–1082. MR 2735011
35. ———, *Well-posedness and existence of bound states for a coupled Schrödinger-gKdV system*, Nonlinear Anal. **73** (2010), no. 8, 2686–2698. MR 2674102
36. M. Essman and J.-P. Shi, *Bifurcation diagrams of coupled Schrödinger equations*, Appl. Math. Comput. **219** (2012), no. 8, 3646–3654. MR 2996805
37. B. Gidas, W.-M. Ni, and L. Nirenberg, *Symmetry and related properties via the maximum principle*, Comm. Math. Phys. **68** (1979), no. 3, 209–243. MR 544879 (80h:35043)

38. ———, *Symmetry of positive solutions of nonlinear elliptic equations in \mathbf{R}^n* , Mathematical analysis and applications, Part A, Adv. in Math. Suppl. Stud., vol. 7, Academic Press, New York, 1981, pp. 369–402. MR 634248 (84a:35083)
39. J. Hulshof and R. van der Vorst, *Differential systems with strongly indefinite variational structure*, J. Funct. Anal. **114** (1993), no. 1, 32–58. MR 1220982 (94g:35073)
40. N. Ikoma, *Uniqueness of positive solutions for a nonlinear elliptic system*, NoDEA Nonlinear Differential Equations Appl. **16** (2009), no. 5, 555–567. MR 2551904
41. P. Korman, *Global solution curves for semilinear systems*, Math. Methods Appl. Sci. **25** (2002), no. 1, 3–20. MR 1874447 (2002k:35114)
42. ———, *Global solution curves for a class of elliptic systems*, Appl. Anal. **89** (2010), no. 6, 849–860. MR 2666544 (2011e:35087)
43. P. Korman and J.-P. Shi, *On Lane-Emden type systems*, Discrete Contin. Dyn. Syst. (2005), no. suppl., 510–517. MR 2192709 (2006h:35061)
44. M.-K. Kwong, *Uniqueness of positive solutions of $\Delta u - u + u^p = 0$ in \mathbf{R}^n* , Arch. Rational Mech. Anal. **105** (1989), no. 3, 243–266. MR 969899 (90d:35015)
45. M.-K. Kwong and L.-Q. Zhang, *Uniqueness of the positive solution of $\Delta u + f(u) = 0$ in an annulus*, Differential Integral Equations **4** (1991), no. 3, 583–599. MR 1097920 (92b:35015)
46. C.-S. Lin, W.-M. Ni, and I. Takagi, *Large amplitude stationary solutions to a chemotaxis system*, J. Differential Equations **72** (1988), no. 1, 1–27. MR 929196 (89e:35075)
47. T.-C. Lin and J.-C. Wei, *Ground state of N coupled nonlinear Schrödinger equations in \mathbf{R}^n , $n \leq 3$* , Comm. Math. Phys. **255** (2005), no. 3, 629–653. MR 2135447 (2006g:35044)
48. O. Lopes, *Stability of solitary waves of some coupled systems*, Nonlinearity **19** (2006), no. 1, 95–113. MR 2191621 (2006i:35304)
49. ———, *Uniqueness of a symmetric positive solution to an ODE system*, Electron. J. Differential Equations (2009), No. 162, 8. MR 2578785 (2011a:34027)
50. L. Ma and B.-Y. Liu, *Symmetry results for decay solutions of elliptic systems in the whole space*, Adv. Math. **225** (2010), no. 6, 3052–3063. MR 2729001 (2011m:35078)
51. L. Ma and L. Zhao, *Uniqueness of ground states of some coupled nonlinear Schrödinger systems and their application*, J. Differential Equations **245** (2008), no. 9, 2551–2565. MR 2455776 (2010a:35042)
52. K. McLeod, *Uniqueness of positive radial solutions of $\Delta u + f(u) = 0$ in \mathbf{R}^n . II*, Trans. Amer. Math. Soc. **339** (1993), no. 2, 495–505. MR 1201323 (94b:35105)
53. T.-C. Ouyang and J.-P. Shi, *Exact multiplicity of positive solutions for a class of semilinear problems*, J. Differential Equations **146** (1998), no. 1, 121–156. MR 1625731 (99f:35061)
54. ———, *Exact multiplicity of positive solutions for a class of semilinear problem. II*, J. Differential Equations **158** (1999), no. 1, 94–151. MR 1721723 (2001b:35117)
55. D. Ruiz, *The Schrödinger-Poisson equation under the effect of a nonlinear local term*, J. Funct. Anal. **237** (2006), no. 2, 655–674. MR 2230354 (2007d:35103)
56. J. Serrin and H.-H. Zou, *Existence of positive solutions of the Lane-Emden system*, Atti Sem. Mat. Fis. Univ. Modena **46** (1998), no. suppl., 369–380. MR 1645728 (99j:35055)
57. B. Sirakov, *Notions of sublinearity and superlinearity for nonvariational elliptic systems*, Discrete Contin. Dyn. Syst. **13** (2005), no. 1, 163–174. MR 2128797 (2006a:35066)
58. ———, *Least energy solitary waves for a system of nonlinear Schrödinger equations in \mathbf{R}^n* , Comm. Math. Phys. **271** (2007), no. 1, 199–221. MR 2283958 (2007k:35477)
59. J. Spruck and Y.-S. Yang, *The existence of nontopological solitons in the self-dual Chern-Simons theory*, Comm. Math. Phys. **149** (1992), no. 2, 361–376. MR 1186034 (93i:58034)
60. ———, *Topological solutions in the self-dual Chern-Simons theory: existence and approximation*, Ann. Inst. H. Poincaré Anal. Non Linéaire **12** (1995), no. 1, 75–97. MR 1320569 (96d:58027)
61. G. Sweers, *Strong positivity in $C(\overline{\Omega})$ for elliptic systems*, Math. Z. **209** (1992), no. 2, 251–271. MR 1147817 (92j:35046)
62. M.-X. Tang, *Uniqueness of positive radial solutions for $\Delta u - u + u^p = 0$ on an annulus*, J. Differential Equations **189** (2003), no. 1, 148–160. MR 1968317 (2004d:35083)
63. E. Timmermans, *Phase separation of Bose-Einstein condensates*, Phys. Rev. Lett. **81** (1998), 5718–5721.
64. W. C. Troy, *Symmetry properties in systems of semilinear elliptic equations*, J. Differential Equations **42** (1981), no. 3, 400–413. MR 639230 (83b:35051)

65. J. Wang and J.-P. Shi, *Standing waves of a weakly coupled Schrödinger system with distinct potential functions*, J. Differential Equations **260** (2016), no. 2, 1830–1864. MR 3419747
66. J.-C. Wei and M. Winter, *Strongly interacting bumps for the Schrödinger-Newton equations*, J. Math. Phys. **50** (2009), no. 1, 012905, 22. MR 2492602 (2010h:35016)
67. J.-C. Wei and W. Yao, *Uniqueness of positive solutions to some coupled nonlinear Schrödinger equations*, Commun. Pure Appl. Anal. **11** (2012), no. 3, 1003–1011. MR 2968605
68. Y.-S. Yang, *Solitons in field theory and nonlinear analysis*, Springer Monographs in Mathematics, Springer-Verlag, New York, 2001. MR 1838682 (2002m:58001)
69. A. C. Yew, A. R. Champneys, and P. J. McKenna, *Multiple solitary waves due to second-harmonic generation in quadratic media*, J. Nonlinear Sci. **9** (1999), no. 1, 33–52. MR 1656377 (99i:78025)
70. L.-G. Zhao, F.-K. Zhao, and J.-P. Shi, *Higher dimensional solitary waves generated by second-harmonic generation in quadratic media*, Calc. Var. Partial Differential Equations **54** (2015), no. 3, 2657–2691. MR 3412388

DEPARTMENT OF MATHEMATICS, SHANGHAI INSTITUTE OF TECHNOLOGY, SHANGHAI, 201418,
P. R. CHINA

E-mail address: `an.yulian@sit.edu.cn`

DEPARTMENT OF MATHEMATICS, NATIONAL CENTRAL UNIVERSITY, CHUNG-LI, TAIWAN 32054,
ROC

E-mail address: `chern@math.ncu.edu.tw`

DEPARTMENT OF MATHEMATICS, COLLEGE OF WILLIAM AND MARY, WILLIAMSBURG, VA 23187-
8795, USA

E-mail address: `jxshix@wm.edu`